

COASSOCIATIVE K3 FIBRATIONS OF COMPACT G_2 -MANIFOLDS

ALEXEI KOVALEV

ABSTRACT. A class of examples of Riemannian metrics with holonomy G_2 on compact 7-manifolds was constructed by the author in [Ko] using a certain ‘generalized connected sum’ of two asymptotically cylindrical manifolds with holonomy $SU(3)$. We consider, on each of the two initial $SU(3)$ -manifolds, a fibration arising from a Lefschetz pencil of K3 surfaces. The gluing of two such K3 fibrations yields a coassociative fibration of the connected sum G_2 -manifold over a 3-dimensional sphere. The singular fibres of this fibration are diffeomorphic to K3 orbifolds with ordinary double points and are parameterized by a Hopf-type link in S^3 . We believe that these are the first examples of fibrations of compact manifolds of holonomy G_2 by coassociative minimal submanifolds.

INTRODUCTION

It is well-known that if a K3 surface contains an elliptic curve then this elliptic curve has self-intersection zero and is a fibre of a holomorphic fibration of the K3 surface over the Riemann sphere. The self-intersection number is calculated by the adjunction formula and the elliptic fibration is defined by a pencil containing the elliptic curve (see [Ha, Propn. 11.1]). All but finitely many fibres of this fibration are smooth and are minimal submanifolds with respect to a Kähler metric on the ambient K3 surface. Moreover, by the Wirtinger theorem the fibres are volume-minimizing in their homology class. Harvey and Lawson [HL] generalized the volume-minimizing property of complex submanifolds of Kähler manifolds by developing a concept of calibrated minimal submanifolds of Riemannian manifolds. One type of examples of calibrated submanifolds found in [HL] is the coassociative submanifolds. These are four-dimensional submanifolds which occur in the Euclidean $\mathbb{R}^7$ and, more generally, in 7-dimensional Riemannian manifolds with holonomy contained in the Lie group G_2 (see §1 for precise definitions).

McLean [McL] studied the deformation theory for calibrated submanifolds. In particular, he showed that local deformations of a smooth compact coassociative submanifold X are always unobstructed and the corresponding ‘moduli space’ is a smooth manifold of dimension $b^+(X)$, the positive index of the cup-product on $H^2(X)$. (This corresponds to the negative index in the actual statement of [McL, Theorem 4.5] as we use a different sign convention for the G_2 -structures.) If $b^+(X) = 3$ then, under some reasonable additional assumptions, the local coassociative deformations of X give a local foliation. This naturally leads to a conjecture that some compact 7-manifolds with holonomy G_2 are fibred by coassociative submanifolds, possibly with singular fibres. Joyce [Jo1, Jo2] constructed first examples of compact 7-manifolds with holonomy G_2 , or G_2 -manifolds; his results include examples of compact coassociative calibrated submanifolds obtained by carefully chosen

orientation-reversing involutions. Later the author gave a series of topologically different examples of compact G_2 -manifolds using a ‘generalized connected sum’ construction [Ko], see also [KLe]. It is also shown in [KN] that one of the G_2 -manifolds obtainable as in [KLe] is a deformation of a G_2 -manifold constructed in [Jo1]. The purpose of this paper is to construct coassociative fibrations of the G_2 -manifolds obtained in [Ko, KLe]. A generic fibre of these fibrations is diffeomorphic to a K3 surface. To the author’s knowledge, these are the first examples of fibrations of *compact* smooth manifolds with holonomy G_2 by *coassociative minimal* submanifolds.

Further motivation for this paper comes from similarities between coassociative submanifolds of G_2 -manifolds and special Lagrangian submanifolds of Calabi–Yau manifolds. Special Lagrangian submanifolds are another instance of calibrated submanifolds appearing in [HL] with unobstructed deformation theory established in [McL]. There are further similarities. For both types of compact calibrated submanifolds, the ‘number of moduli’ of the local deformations is given by some Betti number of the submanifold. Special Lagrangian submanifolds attracted much interest in recent years in connection with the SYZ conjecture [SYZ] which explains mirror symmetry between Calabi–Yau threefolds. The SYZ conjecture was motivated by studies in string theory and inspired works on the mirror symmetry for G_2 -manifolds admitting coassociative fibrations [Ac, GYZ]. A part of the SYZ conjecture asserts that some Calabi–Yau threefolds whose complex structures are close to a certain degenerate limit (the so-called large complex structure limit) admit fibrations by special Lagrangian tori. This may be compared to the requirement that in order to construct coassociative fibrations in this paper we assume that the torsion-free G_2 -structure on a compact 7-manifold M corresponds to a point near the boundary of the moduli space for torsion-free G_2 -structures. (This boundary point is represented by a pair of asymptotically cylindrical manifolds used in the construction of M ; this is made precise in [N, § 5].) More explicitly, a connected sum G_2 -manifold should have a ‘sufficiently long neck’ (as measured by the parameter T in the Main Theorem).

On the other hand, there is a difference concerning properties of the singular fibres. It was shown in [Jo3] that generic fibrations of Calabi–Yau threefolds by special Lagrangian tori are piece-wise smooth and only continuous along the loci of the singular fibres which occur in families of codimension *one*. The coassociative fibrations constructed here are smooth and are continuously differentiable (more precisely, $C^{1,\alpha}$) at the singular fibres. The fibres develop singularities modelled on ordinary double points in the holomorphic deformation families of K3 surfaces, and the singular fibres occur in families of codimension *two*, parameterized by a Hopf-type link in S^3 . This type of coassociative fibration survives under small perturbations of the ambient G_2 -structure, defined by closed 3-forms.

The latter property is crucial as our construction of the coassociative fibrations relies on perturbative analysis, in the form of implicit function theorem in Banach spaces. In fact, we build up on the gluing construction of G_2 -metrics in [Ko] and initially construct an ‘approximating fibration’ whose fibres are coassociative with respect to a G_2 -structure with small torsion. Some familiarity with [Ko] is therefore required and we include in §2 a review of the relevant details before stating the main result and explaining the strategy of the proof in §3. The strategy requires, as a preliminary step, an extension of McLean’s

deformation theory to coassociative K3 orbifolds which we carry out in §4. Appropriate deformation and stability result was recently proved by Lotay [Lo3]. With the deformation set-up in place, we proceed to an implicit function argument for coassociative K3 orbifold fibres in §6 and complete the construction in §7 which deals with the smooth fibres.

Acknowledgements. I am grateful to Simon Donaldson, Dominic Joyce and Miles Reid for helpful discussions on aspects of this project. I also thank Dominic Joyce for valuable comments on an earlier version of this paper.

1. G_2 -STRUCTURES AND COASSOCIATIVE SUBMANIFOLDS

We give a short summary of some standard results concerning the Riemannian geometry associated with the group G_2 and coassociative submanifolds associated with the G_2 -structures. The results given in this section are mostly gathered from [Br, HL, Jo2, McL, Sa] and the reader is referred there for further details.

The group G_2 may be defined as the group of automorphisms of the cross-product algebra on $\mathbb{R}^7$ interpreted as the space of pure imaginary octonions. As the octonions form a normed algebra, any automorphism in G_2 necessarily preserves the Euclidean metric and orientation of $\mathbb{R}^7$. The cross-product can be encoded as an anti-symmetric 3-linear form φ_0 on $\mathbb{R}^7$ defined by $\varphi_0(a, b, c) = \langle a \times b, c \rangle$, where $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product. The 3-form φ_0 and its Hodge dual $*\varphi_0$ are explicitly expressed as

$$\begin{aligned}\varphi_0 &= e_{123} + e_{145} + e_{167} + e_{246} - e_{257} - e_{347} - e_{356}, \\ *\varphi_0 &= e_{4567} + e_{2367} + e_{2345} + e_{1357} - e_{1346} - e_{1256} - e_{1247},\end{aligned}\tag{1.1}$$

where $e_{i_1 \dots i_k} = e_{i_1} \wedge \dots \wedge e_{i_k}$ and $e_i \in (\mathbb{R}^7)^*$, $i = 1, \dots, 7$ is the standard orthonormal co-frame. The group G_2 may be equivalently defined as the subgroup of the orientation-preserving linear isomorphisms $GL_+(7, \mathbb{R})$ fixing φ_0 or the subgroup of $GL_+(7, \mathbb{R})$ fixing $*\varphi_0$, in the action on, respectively, 3- or 4-forms. The group G_2 is a compact Lie group of dimension 14 and thus a proper subgroup of $SO(7)$.

Let M be a smooth oriented 7-manifold. Denote by $\Omega_+^3(M)$ the subset of 3-forms φ on M whose value $\varphi(p)$ at any point $p \in M$ can be identified with φ_0 via some orientation-preserving linear isomorphism $T_p M \rightarrow \mathbb{R}^7$. Any $\varphi \in \Omega_+^3(M)$ induces a G_2 -structure on M ; we shall sometimes, slightly inaccurately, say that φ is a G_2 -structure. The $GL(7, \mathbb{R})$ -orbit of φ_0 is open in $\Lambda^3(\mathbb{R}^7)^*$, so the subset $\Omega_+^3(M) \subset \Omega^3(M)$ is open in the uniform norm topology. As $G_2 \subset SO(7)$ any $\varphi \in \Omega_+^3(M)$ induces a metric $g(\varphi)$ and an orientation of M , so that the image of $e_1, \dots, e_7$ via the isomorphisms $T_p M \rightarrow \mathbb{R}^7$ identifying φ_p with φ_0 is an orthonormal positive co-frame at each $p \in M$. The 4-form $*_\varphi \varphi$ on M defined using the Hodge star of the metric $g(\varphi)$ is, in a similar sense, point-wise isomorphic to $*\varphi_0$ in (1.1).

The metric $g(\varphi)$ induced by a G_2 structure $\varphi \in \Omega_+^3(M)$ will have holonomy contained in G_2 if and only if

$$d\varphi = 0, \quad d*_\varphi \varphi = 0,\tag{1.2}$$

the G_2 -structure φ then is said to be *torsion-free*. Conversely, if the holonomy of a metric g is contained in G_2 then there is a $\varphi \in \Omega_+^3(M)$ satisfying (1.2) so that $g = g(\varphi)$. [Sa, Lemma 11.5] The pair (M, φ) then is called a G_2 -manifold and a metric $g(\varphi)$ induced by

a torsion-free G_2 -structure on a 7-manifold M is called a G_2 -metric. On a compact M , the holonomy of a G_2 -metric is exactly G_2 if and only if the fundamental group of M is finite [Jo2, Propn. 10.2.2].

Now we recall the concept of coassociative submanifolds. Any 4-tuple of orthogonal unit vectors $v_1, v_2, v_3, v_4 \in \mathbb{R}^7$ satisfies the inequality

$$*\varphi_0(v_1, v_2, v_3, v_4) \leq 1. \quad (1.3)$$

If the equality is attained in (1.3) then the oriented 4-dimensional subspace of $\mathbb{R}^7$ defined by v_1, v_2, v_3, v_4 is called a *coassociative subspace*. More generally, if M is a 7-manifold and $\varphi \in \Omega_+^3(M)$ is a G_2 -structure form then any oriented 4-dimensional submanifold $X \subset M$ satisfies

$$*_\varphi \varphi|_X \leq \text{vol}_X, \quad (1.4)$$

in the sense that $*_\varphi \varphi|_X = \alpha \text{vol}_X$ for some real constant $\alpha \leq 1$, where vol_X is the volume form of the metric on X induced by the embedding.

Proposition 1.5 (cf. [HL, Cor. IV.1.20]). *For an orientable 4-dimensional submanifold X of a 7-manifold M with a G_2 -structure $\varphi \in \Omega_+^3(M)$, the equality $*_\varphi \varphi|_X = \text{vol}_X$ is attained in (1.4) for some (necessarily unique) orientation of X if and only if $\varphi|_X = 0$.*

If a G_2 -structure on M satisfies $d*_\varphi \varphi = 0$ then the 4-form $*_\varphi \varphi$ is an instance of a calibration, called *coassociative calibration*, and a 4-dimensional submanifold X with $*_\varphi \varphi|_X = \text{vol}_X$ is then said to be calibrated by $*_\varphi \varphi$. Any compact calibrated submanifold minimizes the volume in its homology class, in particular a compact X calibrated by $*_\varphi \varphi$ is a minimal submanifold of the Riemannian manifold $(M, g(\varphi))$.

The G_2 structures that we deal with in this paper will always be defined by *closed* 3-forms. The following terminology will be in use. A 4-dimensional submanifold $X \subset M$ will be called *coassociative*, with respect to a G_2 -structure $\varphi \in \Omega_+^3(M)$, if $\varphi|_X = 0$. If, in addition, this G_2 -structure satisfies $d*_\varphi \varphi = 0$ (and so $*_\varphi \varphi$ is a calibration) then we say that X is a *coassociative calibrated* submanifold.

A remarkable property of compact coassociative submanifolds is that their local deformations are unobstructed, a result due to McLean [McL]. It uses a ‘tubular neighbourhood theorem’ (e.g. [La, Ch. IV]), which may be stated as follows.

Theorem 1.6. *For any embedded submanifold X of a Riemannian manifold (M, g_M) , the exponential map of g_M ,*

$$v(p) \in (N_{X \subset M})_p \subset T_p M \mapsto \exp_{v(p)} \in M, \quad p \in X,$$

induces a diffeomorphism of a neighbourhood

$$\{v(p) \in N_{X/M} : |v| < C_X \min\{\delta(p), |II(p)|\} \text{ if } v(p) \in T_p M, p \in X\}$$

of the zero section of the normal bundle $N_{X/M}$ onto a neighbourhood of X in M . Here $\delta(p)$ denotes the injectivity radius of g_M and $II(p)$ the second fundamental form at p , $C_X > 0$ is a constant depending only on X .

As $\delta(p)$ depends continuously on $p \in X$ it attains a maximum $\delta > 0$ when X is compact. In that case, the inequality in Theorem 1.6 defining a tubular neighbourhood can be simplified to $|v(p)| < \delta$.

Thus any local deformation of compact coassociative X can be written as $X_v = \exp_v(X)$, for some $v \in \Gamma(N_{X \subset M})$ with small $\|v\|_{C^0}$ in the metric induced from M . Suppose that $\varphi|_X = 0$. A local deformation X_v of X will be coassociative if and only if $F(v) = 0$, where

$$F : v \in \Gamma(N_{X \subset M}) \rightarrow F(v) = (\exp_v)^* \varphi \in \Omega^3(X), \quad (1.7)$$

The normal bundle $N_{X \subset M}$ of a coassociative submanifold is canonically isomorphic and isometric to the bundle of self-dual 2-forms $\Lambda^+ T^* X$ via

$$v \in \Gamma(N_{X/M}) \rightarrow v \lrcorner \varphi|_X \in \Omega^+(X). \quad (1.8)$$

(The G_2 structure 3-forms φ used in [McL] differ from (1.1) by the opposite sign, which leads to $\Omega^-(X)$ rather than $\Omega^+(X)$ in the right-hand side of (1.8).) Composing F with the inverse of (1.8), we obtain a map

$$\tilde{F} : \Omega^+(X) \rightarrow \Omega^3(X).$$

McLean proves:

Theorem 1.9 (cf. [McL], §4). *Suppose that a G_2 -structure on a 7-manifold M is given by a closed 3-form and $X \subset M$ is a coassociative submanifold. Then:*

- (a) *the derivative $(d\tilde{F})_0$ is given by the exterior derivative $d : \Omega^+(X) \rightarrow \Omega^3(X)$, and*
- (b) *the image of $\tilde{F}$ consists of exact forms.*

Remarks. The statement of Theorem 1.9 extracts a part of McLean's results which does not require the compactness of X . The result is stated in [McL] for coassociative calibrated submanifolds but it was later observed in [Go] that the condition $d *_\varphi \varphi = 0$ is not used in the proof.

If X is compact then any exact 3-form on a compact oriented Riemannian 4-manifold is the differential of a self-dual form and Theorem 1.9 sets the scene for an application of the implicit function theorem in Banach spaces. It follows that the local deformations of a compact coassociative X form a smooth manifold of dimension equal to the dimension $b^+(X)$ of harmonic self-dual forms on X , see [McL, Theorem 4.5] or [JS, Theorem 2.5].

Bryant [Br] proved that any closed real-analytic oriented Riemannian 4-manifold X with trivial $\Lambda^+ T^* X$ arises as a coassociative calibrated submanifold in some manifold with torsion-free G_2 -structure. If there are three harmonic self-dual forms on X that are linearly independent at every point then Bryant's result produces examples of non-compact (local) G_2 -manifolds foliated by the coassociative deformations of X .

Finally, an application of McLean's theory shows that compact coassociative submanifolds are 'stable' under small deformations of the G_2 structure. The property will be crucial for the main results of this paper, the following theorem provides an introduction.

Theorem 1.10 (cf. [Jo7, Theorem 12.3.6]). *Suppose that $\varphi(s) \in \Omega^3(M)$, $s \in \mathbb{R}$, is a smooth path of closed G_2 -structure forms on M , and X is a compact submanifold of M such that $\varphi(0)|_X = 0$ and the form $\varphi(s)|_X$ is exact for any s . Then there is an $\varepsilon > 0$*

and for each $|s| < \varepsilon$ a section $v(s)$ of $N_{X/M}$ smoothly depending on s , such that $v(0) = 0$ and $\varphi(s)$ vanishes on $\exp_{v(s)}(X)$.

It is worth to point out the different roles of the two equations in (1.2): the second equation ensures that compact coassociative submanifolds are minimal, whereas the first equation ensures that they have a good deformation theory (Theorems 1.9 and 1.10). Moreover, it is noted in [Br, 0.3.3] that the generic G_2 -structure satisfying $d *_\varphi \varphi = 0$ (but not necessarily $d\varphi = 0$) will not admit any coassociative submanifolds.

2. FROM PENCILS OF K3 SURFACES TO THE APPROXIMATING FIBRATIONS

The method of construction of compact irreducible G_2 manifolds that we shall consider was developed in [Ko]. Because some technical details of this construction will be important in what follows we shall review these details here. We shall also deduce some immediate consequences concerning coassociative submanifolds which are not given in [Ko].

Let W be a Ricci-flat Kähler complex threefold and ω the Kähler form on W . If the holonomy of the Kähler metric is contained in $SU(3)$ (which will be the case e.g. if W is simply-connected) then there is a nowhere-vanishing holomorphic $(3, 0)$ -form Ω on W , sometimes called a holomorphic volume form. We shall call the pair (ω, Ω) a *Calabi–Yau structure* on W . A Calabi–Yau structure (ω, Ω) induces a torsion-free G_2 -structure on the 7-manifold $W \times S^1$ given by the 3-form

$$\varphi_{CY} = \omega \wedge d\theta + \text{Im } \Omega, \quad (2.11)$$

where θ is the standard ‘angle coordinate’ on S^1 (cf. [Jo2, Propn. 11.1.2], our holomorphic volume form differs from the one used there by the factor i). The form φ_{CY} induces the product metric $g_W + d\theta^2$ on $W \times S^1$ corresponding to the Ricci-flat Kähler metric g_W on W and we shall sometimes refer to φ_{CY} as a product G_2 -structure.

If X is a complex surface in W then, for any $\theta_0 \in S^1$,

$$\varphi_{CY}|_{X \times \{\theta_0\}} = \text{Im}(\Omega|_X) = 0,$$

so $X \times \{\theta_0\}$ is a coassociative calibrated submanifold of $W \times S^1$. [McL, Jo2]

Let V be a (non-singular) compact complex threefold with $c_1(V) > 0$, i.e. a Fano threefold, and $D \in |-K_V|$ a K3 surface in the anticanonical linear system of V . Let $D' \in |-K_V|$, $D' \neq D$, be another K3 surface in the anticanonical class, so that $C = D' \cap D$ is a non-singular connected curve in V . Blowing up C we obtain a new threefold $\tilde{V}$ with a holomorphic map $\tau : \tilde{V} \rightarrow \mathbb{CP}^1$ whose fibres are proper transforms of the surfaces in the pencil defined by D and D' .

The proper transform $\tilde{D}$ of D is an anticanonical divisor on $\tilde{V}$ and the complement non-compact complex threefold $W = \tilde{V} \setminus \tilde{D}$ has trivial canonical bundle. We can define a holomorphic coordinate, ζ say, on $\mathbb{CP}^1$, so that $D = \tau^{-1}(0)$. The fibre $\tau^{-1}(\zeta)$ is diffeomorphic to $\tilde{D}$ as a real 4-manifold whenever $|\zeta|$ is sufficiently small, thus the real 6-manifold underlying W has a cylindrical end $\tau^{-1}(\{0 < |\zeta| < \varepsilon\})$ diffeomorphic to $\mathbb{R}_{>0} \times S^1 \times D$. Denote the real coordinates on the first two factors by t, θ and then $\zeta = e^{-t-i\theta}$. The complex structure on the end of W is asymptotic, as $t \rightarrow \infty$, to the product complex

structure on $\mathbb{R}_{>0} \times S^1 \times D$, where D is considered with the complex structure and Kähler form induced by the embedding in V . Note that the latter condition on the Kähler form is *not* restrictive: every Kähler metric on D is obtainable as a restriction of some Kähler metric on V . [Ko] [KLe]

Another class of complex threefolds with similar properties was recently constructed in [KLe]. It uses K3 surfaces with non-symplectic involution.

Recall that by Yau's solution of the Calabi conjecture [Y] the Kähler K3 surface $D \subset V$ has a unique Ricci-flat Kähler metric in its Kähler class. We shall write κ_I for the Kähler form of this metric and $\kappa_J + i\kappa_K$ for a holomorphic volume form on D . The following 'non-compact version of the Calabi conjecture' for W is proved in [Ko] using the results of [TY].

Theorem 2.12 ([Ko], §§ 2–3, 6). *The threefold W is simply-connected and has a complete Ricci-flat Kähler metric g_W of holonomy $SU(3)$. This metric is asymptotically cylindrical in the sense that on the cylindrical end $\tau^{-1}\{0 < |\zeta| < \varepsilon\} \subset W$ the Kähler form ω of g_W has an asymptotic expression*

$$\omega|_{\tau^{-1}\{0 < |\zeta| < \varepsilon\}} = dt \wedge d\theta + \kappa_I + d\psi,$$

and there is a holomorphic volume form Ω on W with an asymptotic expression

$$\Omega|_{\tau^{-1}\{0 < |\zeta| < \varepsilon\}} = (dt + id\theta) \wedge (\kappa_J + i\kappa_K) + d\Psi,$$

where $\zeta = e^{-t-i\theta}$ and the differential forms ψ, Ψ and all their derivatives decay at the rate $O(e^{-\lambda_W t})$ along the end of W , with the exponent $\lambda_W > 0$ depending only on the Kähler metric κ_I on D .

For any W satisfying the assertion of Theorem 2.12, the 7-manifold $W \times S^1$ has a cylindrical end $\mathbb{R}_{>0} \times S^1 \times D \times S^1$ and the product G_2 -structure φ_{CY} on $W \times S^1$ has an asymptotic expression

$$\varphi_{CY}|_{\tau^{-1}\{0 < |\zeta| < \varepsilon\} \times S^1} = \varphi_D + d\tilde{\psi},$$

where

$$\begin{aligned} \varphi_D &= dt \wedge d\theta \wedge d\theta' + \kappa_I \wedge d\theta' + \kappa_J \wedge d\theta + \kappa_K \wedge dt, \\ \tilde{\psi} &= \psi \wedge d\theta' + \text{Im } \Psi, \end{aligned} \tag{2.13}$$

and $d\theta'$ denotes the standard non-vanishing 1-form on the last S^1 factor. Note that the asymptotic model φ_D is determined by the choice of the K3 divisor $D \in |-K_V|$ alone and does not depend on the choice of D' and the resulting pencil on V .

Proposition 2.14. *For a generic choice of Fano threefold V in its deformation family and a generic choice of $D' \in |-K_V|$, the fibres of the map $\tau : \tilde{V} \rightarrow \mathbb{CP}^1$ define a 'generic Lefschetz fibration' in the following sense:*

- (1) *the critical points of τ are non-degenerate (Morse points): if $d\tau(w) = 0$ then the Hessian of τ at w is non-singular, and*
- (2) *any fibre $\tau^{-1}(\zeta)$ contains at most one critical point of τ .*

Proof. This is an application of [Ko] and some standard results in algebraic geometry and we only give an outline of the proof. A generic anticanonical divisor on V is a smooth surface of type K3 [Sh]. The K3 surfaces arising as smooth anticanonical divisors on the deformations of V form a Zariski open subset in a moduli space $\mathcal{K}$ of K3 surfaces whose Picard lattice contains a copy of $H^2(V, \mathbb{Z})$ as a sublattice [Ko, §7]. The moduli space $\mathcal{K}$ is a quasiprojective complex algebraic variety of dimension $20 - b^2(V)$. The degenerations of the K3 surfaces in $\mathcal{K}$ developing an ordinary double point are generic. The anticanonical linear system $|-K_V|$ is parameterized by $\mathbb{C}P^N$ and an application of the Riemann–Roch theorem gives the dimension $N = -K_V^3/2 + 2$, so $N \geq 3$ [Is, Propn. 1.3].

The singular anticanonical divisors on V are therefore parameterized by an algebraic subvariety S of $\mathbb{C}P^N$ of codimension at least 1 and one can show that for a generic Fano threefold in the deformation family of V any connected component of S contains a K3 orbifold with the only singularity an ordinary double point. Each of the conditions (1) and (2) in Proposition 2.14 is an open condition in the Zariski topology of S . Violation of (1) or (2) defines a further subvariety of positive codimension in S . This latter subvariety therefore has codimension at least 2 in $\mathbb{C}P^N$ and can be avoided in the pencil through D and D' with a generic choice of D' . $\square$

If $w_0 \in W$ is a critical point of τ and the Hessian of τ at w_0 is non-degenerate then by the Morse lemma there is a system of local holomorphic coordinates z_j near w_0 , such that $\tau = z_1^2 + z_2^2 + z_3^2$ in these coordinates. The fibre X_0 through w_0 is an orbifold with an isolated singularity $z_1^2 + z_2^2 + z_3^2 = 0$ at w_0 . (For a general theory of orbifolds see [B].) A neighbourhood of w_0 in X_0 admits a local parameterization by uniformizing coordinates $u_1, u_2 \in \mathbb{C}^2/\pm 1$,

$$(u_1, u_2) \mapsto (i(u_1^2 + u_2^2), u_1^2 - u_2^2, 2u_1u_2), \quad (2.15)$$

so this neighbourhood is homeomorphic to a cone on $\mathbb{R}P^3$.

A K3 fibration τ has only finitely many singular fibres. Every singular fibre X_0 of a generic τ (in the sense of Proposition 2.14) is an ‘orbifold K3 surface’ with unique singularity which is an ordinary double point and τ near X_0 defines a one-parameter family of non-singular deformations of X_0 . On the other hand, it is well-known (e.g. [As, §2.6]) that by *blowing up* w_0 one achieves a resolution of the ± 1 orbifold singularity of X_0 which is again a K3 surface (in general, not isomorphic to any nearby fibre of τ) and the exceptional divisor is a complex curve with self-intersection -2 .

The number μ_V of singular fibres of a generic τ is the number of K3 orbifolds in the corresponding generic Lefschetz pencil on V . This number is calculated by an application of Lefschetz theory of hyperplane sections to the topology of algebraic varieties [AF, §5],

$$\mu_V = 2\chi(D) - \chi(C) - \chi(V) = 48 + (-K_V^3) - \chi(V) \quad (2.16)$$

where C is the blow-up locus (the axis of the pencil of K3 surfaces in V), so $-\chi(C) = -K_V^3$ for a Fano threefold V [Is, Propn. 1.6].

We may assume, rescaling ζ by a non-zero constant factor if necessary, that there are no singular fibres of τ on the cylindrical end $\{t > 0\} \subset W$, where $t = -\log|z|$ as before. Extend t to a smooth function, still denoted by t , defined on all of W with $t < 0$ away from

the cylindrical end. Fix once and for all a smooth cut-off function $\alpha(s)$ with $\alpha(s) = 0$, for $s \leq 0$, and $\alpha(s) = 1$, for $s \geq 1$. The 3-form

$$\varphi_{W,T} = \varphi_{CY} - d(\alpha(t - T)\tilde{\psi}) \quad (2.17)$$

defines a G_2 -structure on $W_T \times S^1$ for every sufficiently large T . It interpolates between the product torsion-free G_2 -structure φ_{CY} on $W \times S^1$ induced by the Ricci-flat Kähler structure on W and the product torsion-free G_2 -structure φ_D on the half-cylinder $[T - 1, \infty) \times S^1 \times D \times S^1$ induced by the Calabi–Yau (hyper-Kähler) structure forms $\kappa_I, \kappa_J, \kappa_K$ on D as in (2.13).

Proposition 2.18. *For any $T > 0$, the 3-form φ_T vanishes on each fibre of the map $\tau \times \text{id}_{S^1} : W \times S^1 \rightarrow \mathbb{C}P^1 \times S^1$.*

Proof. It is clear that the claim is true away from the cut-off region $R = [T - 1, T] \times S^1 \times D \times S^1$ because the map τ is holomorphic both in the complex structure of W and in the product complex structure of $\mathbb{R} \times S^1 \times D$ on the end of W .

On the cut-off region, we have

$$\varphi_T|_R = (1 - \alpha)\varphi_{CY} + \alpha\varphi_D - \alpha' dt \wedge \tilde{\psi}$$

and the claim follows as any fibre of τ on the end of $W \times S^1$ is contained in a level set $\{t = \text{const}\}$. $\square$

Now let W_1 and W_2 be two asymptotically cylindrical Calabi–Yau threefolds given by Theorem 2.12 and define $W_j(T) = W_j \setminus \{t_j > T\}$, for any $T > 2$, $j = 1, 2$. Assume that the respective two hyper-Kähler K3 surfaces D_j are ‘hyper-Kähler rotations’ of each other which means that there is an isometry $f : D_1 \rightarrow D_2$ of the Riemannian 4-manifolds such that f^* interchanges the Kähler and holomorphic volume forms of D_1 and D_2 as follows: $\kappa_I^{(2)} \mapsto \kappa_J^{(1)}$, $\kappa_J^{(2)} \mapsto \kappa_I^{(1)}$, $\kappa_K^{(2)} \mapsto -\kappa_K^{(1)}$. (Such an isometry f always exists after some deformations of the Fano threefolds V_1, V_2 used in the construction of W_1, W_2 [Ko, Theorem 6.44]. See also [KLe, Theorem 5.3])

Construct a compact 7-manifold M by joining $W_1(T)$ and $W_2(T)$

$$M = (W_1(T) \times S^1) \cup_{\Upsilon} (W_2(T) \times S^1), \quad (2.19a)$$

identifying collar neighbourhoods of the boundaries via the orientation-preserving diffeomorphism

$$\begin{aligned} \Upsilon : (y, \theta_1, \theta_2, T + t) \in D_1 \times S^1 \times S^1 \times]T + 1, T + 2[\rightarrow \\ (f(y), \theta_2, \theta_1, T + 3 - t) \in D_2 \times S^1 \times S^1 \times]T + 1, T + 2[\end{aligned} \quad (2.19b)$$

Then $\Upsilon^*\varphi_{D_2} = \varphi_{D_1}$, so the two G_2 -structures $\varphi_{j,T} \in \Omega_+^3(W_j(T) \times S^1)$ defined by (2.17) agree on the overlap and together give a well-defined G_2 -structure $\varphi_T \in \Omega_+^3(M)$ on the compact 7-manifold. This latter G_2 -structure form satisfies

$$d\varphi_T = 0, \quad \|d *_{\varphi_T} \varphi_T\|_{L_k^p} < C_{p,k} e^{-\lambda T}, \quad (2.20)$$

for each $T > 2$, where $\lambda < \lambda_{W_j}$, $j = 1, 2$, and λ_{W_j} are defined in Theorem 2.12.

Theorem 2.21 ([Ko], §5). *There exists T_0 and for each $T > T_0$ a 2-form $\eta = \eta_T$ on the 7-manifold M satisfying $\|\eta_T\|_{L_k^p} < K_{p,k}e^{-\lambda T}$ and such that*

$$d * (\varphi_T + d\eta_T) = 0,$$

with the Hodge star $$ in the above formula defined by the metric $g(\varphi_T + d\eta_T)$. Thus the 3-form $\varphi_T + d\eta_T$ induces a torsion-free G_2 -structure on M and the holonomy group of the metric $g(\varphi_T + d\eta_T)$ is G_2 .*

The last claim is true because M is simply-connected. The 3-form φ_T may be thought of as an approximation, improving as $T \rightarrow \infty$, of a torsion-free G_2 -structure on M .

Return, for the moment, to the G_2 -structures $\varphi_{j,T}$ on the two pieces of M . It is clear from the construction (2.19) of M that the fibrations $\tau^{(j)}$ of $W_j(T) \times S^1$ agree on the overlap and hence can be patched to define, for any $T > 1$ a fibration τ_T of M as shown in the following commutative diagram

$$\begin{array}{ccc} W_{1,T} \times S^1 \cup W_{2,T} \times S^1 & \xlongequal{\quad} & M \\ \tau^{(1)} \downarrow & & \tau^{(2)} \downarrow \qquad \qquad \downarrow \tau_T \\ \Delta \times S^1 \cup \Delta \times S^1 & \xlongequal{\quad} & S^3. \end{array} \quad (2.22)$$

where the bottom row is a well-known splitting of the 3-sphere into two solid tori (Δ denotes a disc in $\mathbb{C}P^1$). We obtain from (2.22) and Proposition 2.18 the main result of this section.

Theorem 2.23. *Let (M, φ_T) be a compact 7-manifold with a G_2 -structure constructed from a pair of Fano threefolds, as defined above. Then $\tau_T : M \rightarrow S^3$ is a coassociative fibration defined by (2.22), with respect to φ_T . The singular fibres of τ_T form a subset of codimension 2 in M and are projected by τ onto a link in S^3 .*

This link consists of μ_{V_1} (disjoint) circles in $S^3 \setminus \tau_T(W_1(T) \times S^1)$ and μ_{V_2} (disjoint) circles in $S^3 \setminus \tau_T(W_2(T) \times S^1)$, where μ_{V_j} is defined in (2.16). Each circle in $S^3 \setminus \tau_T(W_j(T) \times S^1)$ is linked, with linking number 1, with each circle in the other subset $S^3 \setminus \tau_T(W_{3-j}(T) \times S^1)$ and is not linked with any circle in $S^3 \setminus \tau_T(W_j(T) \times S^1)$.

Of course, the 4-form $*_{\varphi_T}\varphi_T$ is not in general a calibration on M and the fibres of τ_T are not necessarily calibrated by $*_{\varphi_T+d\eta_T}(\varphi_T + d\eta_T)$. The estimate on the 2-form η_T given in Theorem 2.21 yields an upper bound on $d\eta_T|_X$, for each non-singular fibre X of τ_T ,

$$\|d\eta_T|_X\|_{L_k^p} < \tilde{K}_{p,k}(X)e^{-\lambda T}, \quad (2.24)$$

for each $T > T_0$. The constant $\tilde{K}_{p,k}(X)$ in (2.20) depends on a particular choice of norm (more precisely, on the value of $k - 4/p$) as well as on the choice of X . We shall return to this later, in Proposition 7.64.

3. THE GLUING THEOREM FOR COASSOCIATIVE K3 FIBRATIONS

The estimate (2.20) on the failure of the fibres of τ_T to be calibrated by the 4-form $*_{\varphi_T+d\eta_T}(\varphi_T + d\eta_T)$ also suggests that, the map τ_T might be in some sense an approximation

of a fibration of the holonomy- G_2 manifold $(M, g(\varphi_T + d\eta_T))$ with coassociative calibrated fibres. The next theorem—which is the main result of this paper—asserts that this is indeed the case.

Main Theorem. *Let $(M, g(\varphi_T + d\eta_T))$ be a compact 7-manifold with holonomy G_2 constructed from a pair of Fano threefolds as defined in the previous section and let τ_T be the fibration of M given by Theorem 2.23. There exists, for every sufficiently large T , a diffeomorphism h_T of M onto itself, exponentially close to id_M ,*

$$\sup_{x \in M} \text{dist}_{g(\varphi_T)}(h_T(x), x) < \text{const} \cdot e^{-\lambda T}$$

*with $\lambda > 0$ as determined in Theorem 2.21, and such that the fibres of $\tau_T \circ h_T : M \rightarrow S^3$ are coassociative calibrated by $*_{\varphi_T + d\eta_T}(\varphi_T + d\eta_T)$. In particular, smooth fibres of $\tau_T \circ h_T$ are minimal submanifolds of $(M, g(\varphi_T + d\eta_T))$. The map h_T^{-1} can be taken to be C^1 on the locus of the singular fibres of τ and C^∞ elsewhere on M .*

A generic fibre of $\tau_T \circ h_T$ is diffeomorphic to the real 4-manifold underlying a K3 surface. Each singular fibre is an orbifold diffeomorphic to a ‘K3 orbifold surface’ with one ordinary double point and no other singularities. The discriminant locus (image of the singular fibres) of $\tau_T \circ h_T$ is a link in S^3 , as described in Theorem 2.23.

Remarks. (1) The smooth K3 fibres of $\tau_T \circ h_T$ have $b^+ = 3$ and thus form a maximal deformation family, by McLean’s results [McL]. We shall see, after some additional work below, that the singular fibres of $\tau_T \circ h_T$ also form a maximal deformation family.

Another well-known compact 4-manifold with $b^+ = 3$ is a 4-torus. A fibration by coassociative 4-tori was obtained by Goldstein [Go] for a compact 7-manifold constructed in [Jo1]. However, the G_2 -structure used in [Go] only has a closed 3-form but does *not* define a coassociative calibration (it is close to a calibration and one can obtain an estimate similar to (2.20)), so the fibres need not be minimal submanifolds. The singular coassociative fibres in [Go] have non-isolated singularities and it appears that the problem of perturbing into a map with calibrated fibres would require a different analytic technique than that developed for the fibration (2.22). We hope to return to this problem in a future paper.

(2) A K3 fibre X of τ has trivial normal bundle in M and hence trivial bundle of self-dual forms $\Lambda^+ T^* X$. A trivialization of $\Lambda^+ T^* X$ induces an $Sp(1)$ -structure on X and hence a triple of orthogonal almost complex structures I, J, K (relative to the metric on X) satisfying the quaternionic relations $IJ = -JI = K$. From the construction of the holonomy- G_2 metric on M one can see that one of I, J, K is a deformation of the complex structure induced by embedding of X in the Calabi–Yau threefold W_j . As T tends to infinity, the metric induced by $g(\varphi_T)$ on each fibre of τ_T converges uniformly with all derivatives to a Kähler metric. For fibres near the middle of the neck of M , the limit metric is, moreover, hyper-Kähler. However, there is no general reason for a holonomy reduction for the induced metric on the fibres, for any finite T .

In the rest of the paper we prove the Main Theorem.

We shall construct h_T in the form $h_T = \exp_{v_T}$, for a smooth vector field v_T on M (more precisely, h_T will be obtained as a composition of two exponential maps). A map $\exp_v$ of

the compact Riemannian manifold $(M, g(\varphi_T))$ is well-defined whenever the uniform norm of v is less than the injectivity radius of M . Furthermore, $\exp_v$ defines a *diffeomorphism* of M whenever the uniform norm of both v and its first derivatives is sufficiently small, so that $\exp_v$ is a local diffeomorphism near every point and a bijection of M .

Proposition 3.25. *Let M be a compact 7-manifold with a one-parameter family of Riemannian metrics induced by the G_2 -structures $\varphi_T \in \Omega_+^3(M)$ defined by the connected sum construction, as in §2.*

Then there exists $\varepsilon > 0$ so that for any metric $g(\varphi_T)$, $T \geq 1$, the map $\exp_v$ is a diffeomorphism of M onto itself whenever $\|v\|_{C^1} < \varepsilon$.

Proof. The upper bound ε can be determined by considering restrictions of the metric $g(\varphi_T)$ to small neighbourhoods and taking the supremum. The asymptotically cylindrical properties of the metrics constructed on $W_j \times S^1$ imply that the ε can be taken positive on these manifolds and hence also on M as $g(\varphi_T)$ is exponentially close to the asymptotically cylindrical metrics as $T \rightarrow \infty$. $\square$

We require a C^1 -small vector field $v = v_T$ on M satisfying

$$\exp_v^*(\varphi_T + d\eta_T)|_X = 0$$

for each fibre X of τ_T , given that $\varphi_T|_X = 0$ and η_T can be taken as small as we like by choosing a large T . The construction of such v can be thought of as an infinite-dimensional version of the implicit function problem $F_X(v, s) = 0$ on each fibre X , for a family of vector fields $v = v(s)$, $0 \leq s \leq 1$, with $v(0) = 0$, where

$$F : (v, s) \in \Gamma(TM|_X) \times [0, 1] \rightarrow \exp_v^*(\varphi + s d\eta)|_X \in \Omega^3(X)$$

is a variant of the map appearing in McLean's theory (cf. §1). Here we temporarily dropped the dependence on T from the notation.

If all the coassociative fibres of τ were smooth, then the desired vector field v would be easily obtained by a slight modification of Theorem 1.10 for local deformation families of coassociative submanifolds and then patching together finitely many of these local families, using the compactness of M . However, the fibres of τ develop singularities. Recall also that the deformation problem for a coassociative submanifold is expressed as an equation for sections of vector bundles on the actual submanifold. In light of this, the proof of the Main Theorem naturally falls into two parts concerned, respectively, with the singular fibres of τ_T and the nearby smooth fibres with 'large' curvature.

In order to implement the implicit function strategy we require, in the first place, an extension of the deformation theory to the coassociative *K3 orbifolds* arising as the singular fibres. More precisely, the required property concerns the linearization $(D_1 F)_0$ of the deformation map in v at $v = 0$. This map should be a surjective Fredholm map between appropriate Banach spaces (weighted Sobolev spaces in Theorem 4.35), so the deformations of coassociative K3 orbifold fibres of τ_T are *unobstructed*.

The perturbation h_T of τ_T will be obtained as a composition of exponential maps via the following two results. The proof of Theorem A requires a 'stability' result for coassociative cones defined by complex 2-dimensional cones in $\mathbb{C}^3$. Appropriate stability result for the

tangent cones at the singular points of K3 orbifold fibres of τ_T indeed holds and has been proved by Lotay in [Lo3].

Theorem A. (compare [Lo3]) *Let M be a compact 7-manifold with a smooth one-parameter family of G_2 -structures given by closed 3-forms $\varphi_T \in \Omega_+^3(M)$, $T > T_0$, defined by the generalized connected sum construction in §2. Let $\tau_T : M \rightarrow S^3$, be a coassociative K3 fibration, with respect to φ_T , defined in Theorem 2.23. Suppose that $\varphi_T + d\eta_T$, is a smooth family of torsion-free G_2 -structures on M , such that $\|\eta_T\|_{L_k^p} < K_{p,k}e^{-\lambda T}$ for each $p > 1$, $k \geq 0$.*

Then there exists T_1 and for any $T > T_1$ and $0 \leq s \leq 1$ a $C^{1\alpha}$ vector field $v_{T,s}$ on M , smooth away from the singular fibres of τ_T and satisfying $\|v_{T,s}\|_{L_k^p} < K_{p,k}se^{-\lambda T}$ and $\|v_{T,s}\|_{C^1} < Kse^{-\lambda T}$, with support of v contained in a neighbourhood U of the singular fibres of τ_T , and such that $\varphi + s d\eta_T$ vanishes on every singular fibre of the perturbed fibration $\tau_T \circ \exp_{v_{T,s}}^{-1} : M \rightarrow S^3$. The neighbourhood U may be chosen not to meet the neck of M , i.e. $U \subset (W_1(0) \times S^1 \sqcup W_2(0) \times S^1)$.

We explain in §4 a framework of appropriate weighted Sobolev spaces making $(D_1F)_0$ into a Fredholm map and in §5 show how the surjectivity of $(D_1F)_0$ then follows from Lotay's *stability* result, in the case of conical singularities of K3 orbifold fibres. The surjectivity implies that the deformations of coassociative K3 orbifold fibres of τ_T are *unobstructed*.

The hypothesis of the next result assumes the assertion of Theorem A. We may now deal with an approximating fibration whose singular fibres are precisely coassociative with respect to a G_2 -structure φ_T , for each T .

Theorem B. *Let M be a compact 7-manifold with a smooth one-parameter family of G_2 -structures given by closed 3-forms $\varphi_T \in \Omega_+^3(M)$, $T > T_1$, defined by the generalized connected sum construction in §2. Let $\tau_T : M \rightarrow S^3$ for $T > T_1$ be a coassociative K3 fibration map, with respect to φ_T , defined in Theorem 2.23.*

Then there exists $\varepsilon > 0$ so that if $T > T_1$ and an exact form $d\eta \in \Omega^3(M)$ vanishes on every singular fibre of τ_T and $\|d\eta\|_{C^0(M)} < \varepsilon$, relative to the metric $g(\varphi_T)$, then there is a unique smooth vector field $\tilde{v}_T(\eta)$ on M such that:

(i) $\tilde{v}_T$ vanishes on the singular fibres of τ_T and is point-wise orthogonal to each smooth fibre X of τ_T and $\tilde{v}_T \lrcorner \varphi_T|_X$ is L^2 -orthogonal to the harmonic self-dual forms on X relative to the metric $g(\varphi_T)|_X$;

(ii) $\tilde{v}_T(\eta)$ depends smoothly on T and $d\eta$ and $\|\tilde{v}_T\|_{C^1} = O(\|d\eta\|_{C^1})$;

(iii) $\varphi_T + d\eta$ vanishes on the fibres of $\tau_T \circ \exp_{\tilde{v}_T(\eta)}^{-1}$.

Theorem B is proved in §7.

The estimates of the vector fields $v_{T,s}$ in Theorem A ensure that $\exp_{v_{T,s}}$ for any large T is a well-defined diffeomorphism of M isotopic to id_M . Denote $\tilde{\varphi}_T = \exp_{v_{T,s}}^*(\varphi_T + d\eta_T)$; then, for $T > T_1$, the form $\tilde{\varphi}_T$ vanishes on the singular fibres of τ_T . The form $\tilde{\varphi}_T$ is in the cohomology class of φ_T and $d\tilde{\eta}_T = \tilde{\varphi}_T - \varphi_T$ tends to zero in C^∞ as $T \rightarrow \infty$. For any large T , Theorem B applies to $d\tilde{\eta}_T$ and gives a second vector field $\tilde{v}(\tilde{\eta}_T)$ on M , so that the diffeomorphism $\exp_{\tilde{v}(\tilde{\eta}_T)}$ of M is well-defined and fixes the singular fibres of τ_T . Then

$\varphi_T + d\eta_T$ vanishes on the fibres of $\tau_T \circ h_T$, where

$$h_T = \exp_{v_{T,1}}^{-1} \circ \exp_{\tilde{v}_{T,s}}^{-1}.$$

Thus the fibres of $\tau_T \circ h_T$ are coassociative calibrated by the 4-form $*_{\varphi_T + d\eta_T} \varphi_T + d\eta_T$ and are minimal submanifolds of the holonomy- G_2 manifold $(M, \varphi_T + d\eta_T)$ as required in the Main Theorem.

4. LINEAR ANALYSIS ON COASSOCIATIVE K3 ORBIFOLDS

Before going to prove Theorem A we need to deal with the analytic issues arising in the deformation theory of the singular, K3 orbifold coassociative fibres of the map (2.22). We begin, in this and the next section, by showing that the infinitesimal coassociative deformations of these K3 orbifold fibres are unobstructed.

Our treatment is similar in spirit to one previously used by Joyce in the series of papers on special Lagrangian submanifolds with conical singularities, including [Jo4, Jo6], in that we apply elliptic theory on non-compact manifolds using weighted Sobolev spaces and extend these by certain finite-dimensional spaces to eliminate the obstruction space. Lotay [Lo1] applied the method of [Jo4, Jo6] and other papers in the same series to show that deformations of coassociative submanifolds with conical singularities may in general be obstructed. Recently Lotay developed a rather general deformation and stability theory for coassociative submanifolds whose conical singularities arise from complex cones in $\mathbb{C}^3$.

The ordinary double point singularities of complex surfaces in a threefold W are an instance of conical singularities. On the other hand, complex surfaces in W define coassociative submanifolds in $W \times S^1$ with respect to the product G_2 -structure corresponding to a Calabi–Yau structure on W . As we explain below, using the stability result of [Lo3], in the case of ordinary double point singularities it is possible to set up an unobstructed theory.

Throughout this section, we work on a neighbourhood $M_{\text{loc}} = W_{\text{loc}} \times S^1 \subset M$ of a singular fibre, X_0 say, of the map $\tau = \tau_T$ defined in (2.22). Here $W_{\text{loc}} = (\tau^{(j)})^{-1}(U_0) \subset W_j$ $j = 1$ or 2 , and $U_0 \subset \mathbb{C}P^1$ is an open disc, such that $X_0 \subset W_{\text{loc}}$ and W_{loc} contains no other singular fibres of $\tau^{(j)}$. Respectively, $M_{\text{loc}} = \tau^{-1}(U \times S^1)$; mark a point $0 \in S^1$ and identify W_{loc} with $W_{\text{loc}} \times \{0\} \subset M_{\text{loc}}$. (Note that writing τ rather than τ_T is justified here as there are no singular fibres on the neck of M and the restriction of the fibration map to a neighbourhood away from the neck does not depend on T .) We consider the 7-manifold M_{loc} with the product G_2 -structure

$$\varphi_{CY} = \omega \wedge d\theta + \text{Im } \Omega$$

induced by a Calabi–Yau structure (ω, Ω) on W (as before, θ is an ‘angle coordinate’ on S^1), so φ_{CY} vanishes on the smooth part of X_0 . We shall set up a technical framework to deal with the local coassociative deformations of X_0 , extending McLean’s approach.

Recall that X_0 is an orbifold degeneration of K3 surface with one ordinary double point and no other singularities. Denote by w_0 the ordinary double point point of X_0 and by $X'_0 = X_0 \setminus \{w_0\}$ the complement smooth non-compact complex surface. We shall always use on a neighbourhood of w_0 in W the ‘Morse’ local complex coordinates (z_1, z_2, z_3) discussed

in §2; recall that the local expression of $\tau^{(j)}$ in these coordinates is $z_1^2 + z_2^2 + z_3^2$ and a neighbourhood of w_0 in X_0 corresponds to a neighbourhood of 0 in the cone on $\mathbb{R}P^3$

$$C_0 = \{z_1^2 + z_2^2 + z_3^2 = 0\} \subset \mathbb{C}^3. \quad (4.26)$$

We shall sometimes use on a neighbourhood of w_0 in X_0 the uniformizing coordinates $(u_1, u_2) \in \mathbb{C}^2$ defined in §2, so that

$$(z_1, z_2, z_3) = (i(u_1^2 + u_2^2), u_1^2 - u_2^2, 2u_1u_2). \quad (4.27)$$

The Ricci-flat Kähler metric ω on W_{loc} defines, by restriction, an incomplete Kähler metric on X'_0 and which can be written near w_0 in the form

$$dr^2 + r^2 g_3 + O(r^3), \quad \text{as } r \rightarrow 0. \quad (4.28)$$

where r is the geodesic polar radius at w_0 in W , g_3 is some smooth metric on $\mathbb{R}P^3$, and $o(r^3)$ is understood in the sense of the uniform convergence on $\mathbb{R}P^3$ with all derivatives.

Remark. More explicitly, the metric g_3 can be computed by restricting the inner product on the real tangent space $T_{w_0}W$ induced by ω to the intersection of the unit sphere and the tangent cone of X_0 at w_0 . In the uniformizing coordinates $u_1, u_2, 3$, it is obtained by substituting the expressions (4.27) into (the real part of) $h_{i\bar{j}} dz_i d\bar{z}_j$, where $h_{i\bar{j}}$ is the Hermitian inner product defined by ω at w_0 . Note that the Kähler form ω need not be ‘compatible’ with z_j ’s in any special way. In particular, g_3 is need not in general be induced in the standard way from the round metric on S^3 , nor the ‘obvious’ metric on the link of C_0 induced by the Euclidean metric on $\mathbb{C}^3$.

Note also that the expression for the metric (4.28) in the uniformizing coordinates degenerates at w_0 . In particular, the local 2-form $du_1 \wedge du_2$ on X_0 is smooth in the orbifold sense but its point-wise norm relative to the metric $g(\varphi_{CY})$ blows up at w_0 , $|du_1 \wedge du_2| = O(r^{-1})$ as $r \rightarrow 0$.

In what follows, we extend the local coordinate r to a positive smooth function, still denoted by r , defined on all of X'_0 and such that $r > 1$ away from a coordinate neighbourhood of w_0 .

A *local deformation* of X_0 in M_{loc} is defined as $\exp_v(X_0)$, where v is a vector field on a neighbourhood U of X_0 in M_{loc} with a small C^1 -norm, so that $\exp_v : U \rightarrow M_{\text{loc}}$ is a diffeomorphism of U onto its image. We regard two local deformations $\exp_v(X_0)$ and $\exp_{v'}(X_0)$ as equivalent if $\exp_{v'}(X_0)|_{X_0} = \exp_v(X_0)|_{X_0} \circ \Phi_0$, for some diffeomorphism Φ_0 of X_0 onto itself. (Note that any diffeomorphism of the orbifold X_0 necessarily fixes w_0 .)

The follows is a direct corollary of the tubular neighbourhood Theorem 1.6.

Proposition 4.29. *Let $B_x(\rho)$ denote a ball of radius ρ about zero in the fibre of the normal bundle $N_{X'_0/M_{\text{loc}}}$ over x with the inner product induced by the metric on M_{loc} . There exists $\varepsilon > 0$ such that the Riemannian exponential map defines a diffeomorphism of an open neighbourhood $U_\varepsilon = \cup_{x \in X'_0} B_x(\varepsilon r) \subset N_{X'_0/M_{\text{loc}}}$ onto a neighbourhood of X'_0 in M_{loc} , where r is the ‘polar radius-function’ on X_0 defined above.*

It follows that the bundle isometry $N_{X'_0/M_{\text{loc}}} \cong \Lambda^+ T^* X'_0$ (cf. (1.8)) bijectively identifies local deformations of X_0 fixing w_0 defined by vector fields v point-wise orthogonal to X_0

and the forms $\psi \in \Omega^+(X'_0)$ if the uniform norms of $r^{-1}v$ and $r^{-1}\psi$ are less than ε given by Proposition 4.29. For any such $\psi = v \lrcorner \varphi_{CY}|_{X'_0}$, an orbifold $\exp_v(X_0)$ will be coassociative if and only if ω is a zero of ‘McLean’s map’

$$F : \psi = v \lrcorner \varphi_{CY}|_{X'_0} \in \Omega^+(X'_0) \rightarrow \exp_v^* \varphi_{CY}|_{X'_0} \in \Omega^3(X'_0). \quad (4.30)$$

Recall from Proposition 1.9 that the linearization of F at $\psi = 0$ is an overdetermined-elliptic differential operator $(dF)_0 = d : \Omega^+(X_0) \rightarrow \Omega^3(X_0)$ and that the image of F consists of exact 3-forms on X'_0 . In order to apply the implicit function theorem to F we require a choice of Banach space completions for the space of self-dual forms $v \lrcorner \varphi_{CY}$ on X'_0 arising from local deformations of X_0 and also for the space of bounded exact 3-forms on X'_0 , so that the exterior derivative extends to a *surjective* operator between the two Banach spaces.

The Kähler metric induced on X'_0 from W_{loc} does *not* extend to a smooth orbifold metric on X_0 , so trying to work with orbifold versions of e.g. the usual Sobolev spaces on a compact X_0 and is not a very promising way. Instead, we use a ‘conformal blow-up’ of X'_0 at the singular point and apply the elliptic theory for manifolds with asymptotically cylindrical ends from [LM, MP, Me].

The non-compact submanifold X'_0 is diffeomorphic to a smooth 4-manifold with cylindrical end $\mathbb{R}_{>0} \times \mathbb{R}P^3$ which corresponds to a neighbourhood of w_0 via $r = e^{-t}$, where t is the coordinate on the $\mathbb{R}_{>0}$ factor. Restricting to the end of X'_0 we can write

$$dr^2 + r^2 g_3 + o(r^3) = e^{-2t} g_{\text{cyl}} = e^{-2t} (dt^2 + g_3 + o(e^{-t})), \quad \text{as } t \rightarrow \infty, \quad (4.31)$$

which shows that the metric (4.28) induced on X'_0 from M_{loc} is conformally equivalent to an *asymptotically cylindrical* metric g_{cyl} . In particular, the self-dual forms defined by the metrics g_{cyl} and $\omega|_{X'_0}$ are the same.

We shall need exponentially weighted Sobolev spaces on X'_0 . By definition, $e^{-\delta t} L_k^p(X'_0)_{\text{cyl}}$ is the space of functions $e^{-\delta t} f$ such that $f \in L_k^p(X'_0)$ and the norm is defined by $\|e^{-\delta t} f\|_{e^{-\delta t} L_k^p} = \|f\|_{L_k^p}$. Here we used the subscript ‘cyl’ to indicate that the L_k^p norm in the previous sentence is calculated using the metric g_{cyl} . This will be important when we consider the differential forms on X'_0 .

There is a preferred choice of weight $\delta = k - 4/p$. Denote the corresponding weighted spaces by $W_k^p(X_0)$. In terms of the radial parameter r on X'_0 the W_k^p -norm is expressed as

$$\|f\|_{W_k^p(X_0)} = \left\| \frac{f}{r^k} \right\|_p + \left\| \frac{\nabla f}{r^{k-1}} \right\|_p + \dots + \|\nabla^k f\|_p.$$

The above expression for W_k^p norm extends to the differential forms of any degree m on X'_0 but note that the point-wise norms of the m -forms are rescaled by the ‘conformal weight’ factor e^{-mt} when passing to the metric g_{cyl} . In view of this, we define

$$W_k^p \Omega^m(X_0) = e^{-(k-4/p+m)t} L_k^p \Omega^m(X'_0)_{\text{cyl}}.$$

Then the exterior derivative extends to a bounded linear map $W_k^p \Omega^m(X'_0) \rightarrow W_{k-1}^p \Omega^{m+1}(X'_0)$.

There is a simple relation between the W_k^p spaces and the usual, unweighted Sobolev spaces $L_k^p(X_0)$. The following result is proved in [Bi] in the case of a flat Euclidean ball (with r the Euclidean distance to the origin). However, the argument of the proof works, with

only a change of notation, for a punctured ball endowed with a metric having the ‘conical’ form (4.28). Considering a punctured neighbourhood of w_0 in X_0 as the quotient of a 4-dimensional punctured ball with respective ± 1 -invariant metric and restricting attention to ± 1 -invariant functions we obtain.

Proposition 4.32 (cf. [Bi, Theorem 1.3]). *Suppose that ℓ is a non-negative integer such that $\ell - 1 < k - 4/p < \ell$ and let $\delta = k - 4/p$. Then one has*

$$W_k^p(X_0) = \{f \in L_k^p(X_0) \mid \lim_{w \rightarrow w_0} \nabla^m f(w) = 0, \text{ for all } 0 \leq m \leq \ell - 1\} \quad (4.33)$$

and the W_k^p -norm on the left-hand side is equivalent to the L_k^p -norm on the right-hand side.

The vanishing condition in the right-hand side of (4.33) makes sense as L_k^p embeds in $C^{\ell-1}$. Proposition 4.32 extends in the usual way to sections of vector bundles over X_0 by considering a ± 1 -equivariant local trivialization near w_0 .

In view of the local regularity results for coassociative calibrated manifolds [HL, § IV.2.7] we require a Banach space consisting of the C^1 self-dual forms on X'_0 . We shall use the completion of the space of self-dual forms in a W_k^p -norm fixing a choice of $p > 1$, $k \in \mathbb{Z}$ such that

$$1 < k - 4/p < 2 \quad (4.34)$$

With this choice, every self-dual form $\psi \in W_k^p \Omega^+(X'_0)$ vanishes to order two at w_0 , in particular ψ is Lipschitz continuous at w_0 with any Lipschitz constant $\varepsilon > 0$. Whenever the W_k^p norm of ψ is small, the corresponding section v of $N_{X'_0/M_{\text{loc}}}$ defines a local deformation $\exp_v$ of X_0 which fixes the singular point w_0 and the tangent cone at w_0 .

In order to include local deformations of X_0 which move the tangent cone and the singular point, we extend the weighted Sobolev space of self-dual forms by adding a finite-dimensional space

$$E_0 = \{v_{\mathbf{e},L} \lrcorner \varphi_{CY}|_{X'_0} : \mathbf{e} \in T_{w_0} M_{\text{loc}}, L \in \text{End}(T_{w_0} M_{\text{loc}})\}.$$

Here $v_{\mathbf{e},L}$ denotes a choice of a smooth vector field on M_{loc} smoothly depending on the parameters $\mathbf{e}, L$, such that $v_{\mathbf{e},L}(w_0) = \mathbf{e}$ and $dv_{\mathbf{e},L}(w_0) = L$. In the last condition we define $dv_{\mathbf{e},L}$ by using the coordinates z_i, θ near $w_0 \in M_{\text{loc}}$ to express $v_{\mathbf{e},L}$ locally as a smooth map from a neighbourhood of zero in T_{w_0} to T_{w_0} . The Banach space $E_0 + W_k^p \Omega^+(X'_0)$, for $1 < k - 4/p < 2$, does not depend on the choice of vector fields $v_{\mathbf{e},L}$ given above as the ambiguity is $O(r^2)$, $r \rightarrow 0$, which is contained in W_k^p by Proposition 4.32. It is not difficult to check, using Proposition 4.32, that for every smooth local deformation $\exp_v$ of X_0 the self-dual form $v \lrcorner \varphi_{CY}|_{X'_0}$ is in $E_0 + e^{-\delta t} L_k^p \Omega^+(X'_{0,\text{cyl}})$. On the other hand, straightforward calculation in local coordinates on M_{loc} near w_0 shows that any smooth exact 3-form on M_{loc} restricts to a form in $d(E_0) + W_{k-1}^p(\Omega^3(X'_0))$. (As $k-1-4/p > 0$, the space $W_{k-1}^p(\Omega^3(X'_0))$ alone only contains forms vanishing at w_0 .)

We are now ready to state the main result of this section.

Theorem 4.35. *If the Sobolev space parameters satisfy (4.34) then the exterior derivative defines a bounded linear map between Banach spaces*

$$d : E_0 + W_k^p \Omega^+(X'_0) \rightarrow d(E_0) + \{\eta \in W_{k-1}^p \Omega^3(X'_0) : d\eta = 0\} \quad (4.36)$$

which is surjective and has a one-dimensional kernel spanned by the Kähler form $\omega|_{X'_0}$.

Remark. The 1-dimensional kernel of (4.36) arises from the S^1 -symmetry of the torsion free G_2 -structure φ_{CY} on $W \times S^1$. Thus Theorem 4.35 shows that the S^1 -families of coassociative K3 orbifolds arising in the approximating fibration (2.22) are *maximal* deformation families (the smooth fibres, of course, have the same property by Theorem 1.9 and [McL] as $b^+ = 3$ for a K3 surface).

5. PROOF OF THEOREM 4.35

Remark. Putting $x = e^{-t}$ we can think of the manifold X'_0 as the interior of a compact manifold with boundary $\overline{X'_0} = X'_0 \cup (\{x = 0\} \times \mathbb{R}P^3)$ obtained by adding a copy of $\mathbb{R}P^3 = \{x = 0\}$ ‘at infinity’. (The added $\mathbb{R}P^3$ may also be canonically identified with the unit spherical space form $S^3/\pm 1$ in the tangent cone of $X_0 \subset W_{\text{loc}}$ at w_0 .) The asymptotically cylindrical metric g_{cyl} on $\overline{X'_0}$ can be written in the form

$$g_{\text{cyl}} = \frac{dx^2}{x^2} + \tilde{g}, \quad (5.37)$$

where $\tilde{g}$ is a symmetric semi-positive definite form smooth up to the boundary: at any point in $\{x = 0\}$, $\tilde{g}$ is smooth in the $\mathbb{R}P^3$ directions and has one-sided derivatives in x of any order. Then (5.37) gives an instance of a ‘smooth exact b -metric’, as defined by Melrose [Me, Ch. 2]. The results proved in [Me] for manifolds with smooth exact b -metrics can therefore be applied to (X'_0, g_{cyl}) .

It will be convenient to reduce Theorem 4.35 to a result concerning a linear operator between the W_k^p -spaces alone. We look at the kernel first. (The subscript ‘cyl’ at the exponentially weighted spaces on X'_0 will now be dropped from the notation.)

Proposition 5.38. *Suppose that $k - 4/p > 1$. The map $d : e^{-\delta t} L_k^p \Omega^+(X'_0) \rightarrow e^{-\delta t} L_{k-1}^p \Omega^3(X'_0)$ has a 3-dimensional kernel for any $0 \leq \delta < 1$, a 1-dimensional kernel for any $1 \leq \delta < 2$, and is injective for any $\delta \geq 2$.*

Proof. This is an application of the Hodge theory on asymptotically cylindrical manifolds. The kernel of d is contained in the kernel of the Laplacian acting on $e^{-\delta t} L_k^p \Omega^+(X'_0)$. Furthermore, the standard integration by parts argument is valid when $\delta > 0$ and shows that the two kernels coincide. By [APS, Propn. 4.9] or [Me, Propn. 6.14], the L^2 -kernel of the Laplacian on the m -forms on an asymptotically cylindrical manifold X'_0 is isomorphic to the image of the natural inclusion homomorphism $H_c^m(X'_0) \rightarrow H^m(X'_0)$ of the de Rham cohomology groups, where the subscript ‘c’ indicates the cohomology with compact support.

Considering the exact sequence of the de Rham cohomology groups

$$\dots \rightarrow H^{m-1}(\mathbb{R}P^3) \rightarrow H_c^m(X'_0) \rightarrow H^m(X'_0) \rightarrow H^m(\mathbb{R}P^3) \rightarrow \dots$$

we find that $H_c^0(X'_0) = 0$, $H_c^4(X'_0) \cong \mathbb{R}$ and the inclusion homomorphism $H_c^m(X'_0) \rightarrow H^m(X'_0)$ is an isomorphism for $1 \leq m \leq 3$. Recall from §2 that X'_0 is isomorphic to the complement of a (-2) -curve in a K3 surface, X say; this (-2) -curve is topologically a

sphere. Considering the Maier–Vietoris exact sequence for the union of X'_0 and a tubular neighbourhood of the ‘missing’ (-2) -curve we find that $H^2(X'_0)$ is a complement in $H^2(X)$ of the one-dimensional subspace generated by the Poincaré dual of the (-2) -curve. The cup-product on $H_c^2(X'_0)$ has maximal positive subspace $H^+(X'_0)$ of dimension 3 and so the L^2 -kernel of the Laplacian on $\Omega^+(X'_0)$ is 3-dimensional.

It is easy to identify, for $0 < \delta < 1$, the 3-dimensional space of $O(e^{-\delta t})$ exponentially decaying closed self-dual forms on X'_0 . This space is spanned by the restriction $\omega|_{X_0}$ of the Kähler form on W_{loc} and the restrictions of the real and imaginary parts of the holomorphic $(2, 0)$ -form $((\partial\tau)^\sharp \lrcorner \Omega)|_{X_0}$, defined using the K3 fibration map τ on W_{loc} . The Kähler form $\omega|_{X_0}$, measured with the metric g_{cyl} , is $O(e^{-2t})$, as $t \rightarrow \infty$, but not $O(e^{-(2+\varepsilon)t})$ for any $\varepsilon > 0$. As the $(1, 0)$ -form $\partial\tau$ has a zero of order 1 at w_0 , the vector field $(\partial\tau)^\sharp$ is $O(1/r)$, as $r \rightarrow 0$, when measured in the Kähler metric $\omega|_{X_0}$. Hence the real and imaginary parts of $((\partial\tau)^\sharp \lrcorner \Omega)|_{X_0}$ are $O(e^{-t})$ but not $O(e^{-(1+\varepsilon)t})$ in the metric g_{cyl} .

Neither of the three self-dual forms spanning the L^2 -kernel of the Laplacian (and the L^2 -kernel of A) is in $e^{-\delta t}L_k^p$ for $\delta > 2$. $\square$

Corollary 5.39. *If $1 < k - 4/p < 2$ then the map 4.36 has a one-dimensional kernel spanned by the Kähler form $\omega|_{X'_0}$.*

Proof. It is not difficult to check that $\omega|_{X'_0} \in E_0 + W_k^p\Omega^+(X'_0) \subset e^{-\delta t}L_k^p\Omega^+(X'_0)$, for any $1 < \delta < 2$ but $E_0 + W_k^p\Omega^+(X'_0)$. $\square$

The next two lemmas determine the codimensions of the relevant W_k^p spaces.

Lemma 5.40. *If $1 < k - 4/p < 2$ then the codimension of $W_k^p\Omega^+(X'_0)$ in $E_0 + W_k^p\Omega^+(X'_0)$ is 41.*

Proof. If $\mathbf{e} \neq 0$ then $|e^{2t}v_{\mathbf{e}, L \lrcorner \varphi_{CY}}|_{g_{\text{cyl}}}$ has a non-zero lower bound on an open subset of the cylindrical end, of the form $\mathbb{R}_+ \times U'$ where U' is open in $\mathbb{R}^3$. So $v_{\mathbf{e}, L \lrcorner \varphi_{CY}}$ with $\mathbf{e} \neq 0$ is never in $W_k^p\Omega^+(X'_0)$ if $k - 4/p > 0$.

The form $v_{0, L \lrcorner \varphi_{CY}}$ will be in $W_k^p\Omega^+(X'_0)$ with $1 < k - 4/p < 2$ precisely if L leaves invariant the tangent cone of X_0 at w_0 . This, in turn, will be the case if and only if the Zariski tangent space $T_{w_0}W_{\text{loc}}$ of X_0 is an invariant subspace of L and the restriction of L to $T_{w_0}W_{\text{loc}}$ is up to a complex factor an element of $SO(3, \mathbb{C})$ (so that L preserves the tangent cone (4.26)). We find that the subspace of endomorphisms in $\text{End } T_{w_0}M_{\text{loc}}$ preserving the tangent cone has dimension 15. As the dimension of E_0 is $7 + 49$ the result follows. $\square$

Lemma 5.41. *If $1 < k - 4/p < 2$ then the codimension of $W_{k-1}^p\Omega^3(X'_0) \cap \text{Ker } d$ in $d(E_0) + (W_{k-1}^p\Omega^3(X'_0) \cap \text{Ker } d)$ is 18.*

Proof. Recall from §4 that the space $d(E_0) + (W_{k-1}^p\Omega^3(X'_0) \cap \text{Ker } d)$ contains the restrictions to X'_0 of all the smooth exact 3-forms on M_{loc} . On the other hand, $W_{k-1}^p\Omega^3(X'_0)$ is precisely the space of L_{k-1}^p 3-forms on X'_0 with zero limit at w_0 . Since $X_0 \subset W_{\text{loc}}$ and the Zariski tangent space of X_0 at w_0 is $T_{w_0}W_{\text{loc}}$ the codimension of interest may be computed as $\dim \Lambda^3 T_{w_0}^* W_{\text{loc}} - \dim(d(E_0) \cap W_{k-1}^p\Omega^3(X'_0))$.

A calculation in the local complex coordinates on W_{loc} near w_0 shows that $d(v_{\mathbf{e},L} \lrcorner \varphi_{CY})|_{X'_0} \in W_{k-1}^p \Omega^3(X'_0)$ holds precisely if $d(v_{\mathbf{e},L} \lrcorner \varphi_{CY})|_{W_{\text{loc}}}(w_0)$ is the real or imaginary part of a $(3,0)$ -form on $T_{w_0} W_{\text{loc}}$. $\square$

From Corollary 5.39 and Lemmas 5.40, 5.41 and some straightforward linear algebra we find that Theorem 4.35 is equivalent to the following technical result on an asymptotically cylindrical 4-manifold (X'_0, g_{cyl}) .

Theorem 4.35'. *If $1 < k - 4/p < 2$ then the injective linear map*

$$A : \psi \in W_k^p \Omega^+(X'_0) \rightarrow d\psi \in W_{k-1}^p \Omega^3(X'_0) \cap \text{Ker } d, \quad (5.42)$$

has a 22-dimensional cokernel.

In the remainder of this section we prove Theorem 4.35'. Note that the injectivity of A follows from Proposition 5.38. Therefore, the dimension of $\text{Coker } A$ is minus the Fredholm index of A . For the index computation, it is convenient to observe that the map A is equivalent to a component of an elliptic operator

$$D : (f, \psi) \in e^{-\delta t} L_k^p(\Omega^0 \oplus \Omega^+)(X'_0) \rightarrow df - *d\psi \in e^{-\delta t} L_{k-1}^p \Omega^1(X'_0). \quad (5.43)$$

For any real δ , we write $\text{index}_{-\delta} D$ to indicate that D is considered on the $e^{-\delta t}$ -weighted Sobolev spaces. A similar notation will be used for A and for the kernels and cokernels.

Proposition 5.44. $\text{index}_{-\delta} A = \text{index}_{-\delta} D$ for any $\delta > 0$.

Proposition 5.44 will be deduced from the following.

Lemma 5.45. *If $\varepsilon > 0$ is sufficiently small then $\text{Coker}_{-\varepsilon} D = \{0\}$.*

Proof of Proposition 5.44 assuming Lemma 5.45. For any $\delta > 0$, the cohomology of the weighted de Rham complex on X'_0

$$e^{-\delta t} L_k^p \Omega^0(X'_0) \xrightarrow{d} e^{-\delta t} L_{k-1}^p \Omega^1(X'_0) \xrightarrow{d} e^{-\delta t} L_{k-2}^p \Omega^2(X'_0) \xrightarrow{d} \dots \quad (5.46)$$

is isomorphic to the de Rham cohomology with compact support $H_c^*(X'_0)$ [Me, Propn. 6.13]. The formal L^2 -adjoint of (5.46) is a complex of the $e^{\delta t}$ -weighted spaces. The cohomology of the latter complex at the Ω^m -term is isomorphic to the de Rham cohomology $H^{4-m}(X'_0)$.

Recall from the proof of Proposition 5.38 that $H_c^0(X'_0) = H_c^1(X'_0) = 0$. So the exterior derivative maps $e^{-\delta t} L_k^p \Omega^0(X'_0)$, for each $\delta > 0$, isomorphically onto the subspace of closed forms in $e^{-\delta t} L_{k-1}^p \Omega^1(X'_0)$. Standard integration by parts argument of the Hodge theory for compact manifolds is valid for the exponentially decaying forms on X'_0 and shows that the spaces of 1-forms $d(e^{-\delta t} L_k^p \Omega^0(X'_0))$ and $d^*(e^{-\delta t} L_k^p \Omega^+(X'_0))$ are L^2 -orthogonal if $\delta > 0$. By Lemma 5.45, D is a surjective operator between $e^{-\varepsilon t}$ -weighted spaces for any small $\varepsilon > 0$, so we obtain a decomposition

$$e^{-\varepsilon t} L_{k-1}^p \Omega^1(X'_0) = d(e^{-\varepsilon t} L_k^p \Omega^0(X'_0)) \oplus d^*(e^{-\varepsilon t} L_k^p \Omega^+(X'_0)).$$

For an arbitrary $\delta > 0$ we can write any $\xi \in e^{-\delta t} L_{k-1}^p \Omega^1(X'_0)$ as $\xi = d\xi_0 + *d\xi_+$ where $\xi_+ \in e^{-\varepsilon t} L_k^p \Omega^+(X'_0)$ for some *small* $\varepsilon > 0$ but $\xi_0 \in e^{-\delta t} L_k^p \Omega^0(X'_0)$ (as explained above). Therefore, the image of D on the $e^{-\delta t}$ -weighted spaces is complemented by d^* -closed forms and the proposition follows. $\square$

For the proof of Lemma 5.45, we need to recall from [LM] or [Me] some Fredholm theory for elliptic operators on asymptotically cylindrical Riemannian manifolds. Let $\Lambda_\infty^+ T^*(\mathbb{R}_{>0} \times \mathbb{R}P^3)$ denote the bundle of self-dual forms with respect to the metric $dt^3 + g_3$, the asymptotic model of g_{cyl} . A point-wise orthogonal projection, relative to $dt^3 + g_3$, $\sigma : \Lambda^+ T^*(\mathbb{R}_{>0} \times \mathbb{R}P^3) \rightarrow \Lambda_\infty^+ T^*(\mathbb{R}_{>0} \times \mathbb{R}P^3)$ defines a bundle isomorphism asymptotic to the identity as $t \rightarrow \infty$. The coefficients of D are determined by the metric g_{cyl} and D is asymptotic, on the end of X'_0 , to an operator $D_\infty \circ (1 \oplus \sigma)$, where D_∞ is given by the same formula as in (5.43) but using the product cylindrical metric $dt^3 + g_3$ rather than g_{cyl} . The coefficients of the operator D_∞ on $\mathbb{R} \times \mathbb{R}P^3$ are independent of t .

Proposition 5.47. (i) *The elliptic operator D between $e^{-\lambda t}$ -weighted Sobolev spaces, is Fredholm if and only if $d(\lambda) = 0$, where*

$$d(\lambda) = \dim\{e^{-\lambda t}p(t, y) \mid p(t, y) \text{ is polynomial in } t \text{ and } D_\infty(e^{-\lambda t}p(y, t)) = 0\}. \quad (5.48)$$

The set $\{\delta \in \mathbb{R} : d(\delta) \neq 0\}$ is discrete in $\mathbb{R}$.

(ii) *The index of D is independent of p, k but depends on the weight parameter λ according to the formula*

$$\text{index}_{\delta''} D - \text{index}_{\delta'} D = \sum_{\delta' < \lambda < \delta''} d(\lambda) \quad (5.49)$$

for any $\delta' < \delta''$ such that $d(\delta') \neq 0$, $d(\delta'') \neq 0$.

(iii) *The kernel of D consists of smooth forms and is independent of p, k . The cokernel of D can be identified with the kernel of the formal L^2 -adjoint of D with respect to g_{cyl} ,*

$$D^* = d^* \oplus d^+ : e^{\delta t} L_{k-1}^p \Omega^1(X'_0) \rightarrow e^{\delta t} L_k^p (\Omega^0 \oplus \Omega^+)(X'_0).$$

In particular, $\dim \text{Coker}_{-\lambda} D = \dim \text{Ker}_\lambda D^$ if $d(\lambda) = 0$.*

Proposition 5.47 is a direct application of [LM] or [Me]. In the case of D we don't need to worry about the possibility of $\lambda \in \mathbb{C} \setminus \mathbb{R}$ with $d(\lambda) \neq 0$, as we shall see in a moment.

Proof of Lemma 5.45. The operator D_∞ can be expressed, using some vector bundle isomorphisms, as $\partial_t + D^{(3)}$ where $D^{(3)}$ is a formally self-adjoint operator on $\mathbb{R}P^3$ whose square is the Laplacian on $(\Omega^0 \oplus \Omega^1)(\mathbb{R}P^3)$ (cf. e.g. [MMR, pp. 132–134]). This yields $d(0) = 1$. Therefore, D is a Fredholm operator on $e^{-\varepsilon t}$ - and on $e^{\varepsilon t}$ -weighted spaces whenever $\varepsilon > 0$ is sufficiently small. Furthermore, for a small $\varepsilon > 0$, we obtain

$$\dim \text{Ker}_{-\varepsilon} D = 3, \quad \dim \text{Ker}_{-\varepsilon} D^* = 0$$

by the Hodge theory arguments as in the proof of Proposition 5.38, using also the relation $DD^* = dd^* + \frac{1}{2}d^*d$. On the other hand,

$$\text{index}_\varepsilon D - \text{index}_{-\varepsilon} D = 1,$$

by (5.49), whence

$$\dim_{-\varepsilon} \text{Coker } D + \dim_\varepsilon \text{Ker } D = 4.$$

by Proposition 5.47(iii). But $\dim_\varepsilon \text{Ker } D \geq 1 + \dim_{-\varepsilon} \text{Ker } D = 4$ as the $e^{\varepsilon t} L_k^p$ -kernel contains the constants, so $\text{Coker}_{-\varepsilon} D = \{0\}$. $\square$

The next result is proved by Lotay [Lo3] and is essentially the stability property for the coassociative cones defined by complex cones biholomorphic to C_0 . Clause (i) will be needed for the $C^{1,\alpha}$ -regularity claim in the next section.

Proposition 5.50 (J. D. Lotay). *Let D be the elliptic operator defined in (5.43) over a K3 orbifold X_0 and let $d(\lambda)$ be as defined in (5.48). Then*

- (i) *any $0 < \lambda \leq 0$ such that $d(\lambda) \neq 0$ is an integer;*
- (ii) *$\sum_{0 < \lambda < \delta} d(\lambda) = 25$, for some $3 < \delta < 4$.*

Theorem 4.35' follows from Proposition 5.50 by dimension counting. For if $3 < \delta < 4$ then using (5.49), Propositions 5.38 and 5.44, and Proposition 5.50(ii) we obtain $d(\delta) \neq 0$ and

$$\dim \text{Coker}_{-\delta} A = -\text{index}_{-\delta} A = -\text{index}_{-\delta} D = \sum_{0 < \lambda < \delta} d(\lambda) - 3 = 22$$

as required.

Remarks on Proposition 5.50. The conformal factor at the Hodge star in (5.43) is precisely the inverse of the conformal weight of 1-forms. Therefore, the operator D and its asymptotic model D_∞ are *conformally invariant* and can be interchangeably considered on X'_0 with the Kähler metric $\omega|_{X'_0}$.

The asymptotic model for the cylindrical end of X'_0 corresponds to the cone C_0 in $T_{w_0}W_{\text{loc}} \cong \mathbb{C}^3$ defined by (4.26) and the asymptotic model for g_{cyl} is conformally equivalent to the Kähler metric $dr^2 + r^2 g_3$ on C_0 induced by restricting from $T_{w_0}W_{\text{loc}}$ a Hermitian inner product defined by $\omega(w_0)$. The kernel elements of D_∞ contributing to the dimensions $d(\lambda)$ are expressed on C_0 as pairs

$$(r^\lambda p_0(\log r, y), r^{\lambda-2} p_+(\log r, y) \in (\Omega^0 \oplus \Omega^+)(C_0 \setminus \{0\}), \quad (5.51)$$

where a 0-form p_0 and a self-dual form p_+ are polynomial in $\log r$ and smooth in $y \in \mathbb{R}P^3$. As noted earlier, the conformal rescaling produces an extra factor r^{-2} for the self-dual forms in (5.51).

For a Kähler surface $C_0 \setminus \{0\}$, there is a commutative diagram, cf. [FM, §4.3.3],

$$\begin{array}{ccc} \Omega^0 \oplus \Omega^+ & \xrightarrow{D_\infty} & \Omega^1 \\ \iota, j + \pi^{0,2} \downarrow \cong & & \cong \downarrow \pi^{0,1} \\ \Omega^0 \oplus \Omega^{0,2} & \xrightarrow{2\bar{\partial} + \bar{\partial}^*} & \Omega^{0,1}, \end{array} \quad (5.52)$$

where Ω^0 in the bottom row denotes the complex-valued 0-forms, ι is the inclusion map, $j(\alpha) = -i\alpha \cdot \omega_{C_0}$ is the contraction with the Kähler form and the inverses of $\pi^{p,q}$ map the respective complex vector space to its underlying real vector space.

Suppose that

$$2\bar{\partial}f + \bar{\partial}^*\psi = 0 \quad (5.53)$$

and that $(f, \psi) \in \Omega^0 \oplus \Omega^{0,2}$ is obtained from (5.51) for some $\lambda > 0$ via the left column of (5.52). Then since $\text{Re } f$ has degree r^λ , whereas $\text{Im } f$ and ψ both have degree $r^{\lambda-2}$ we deduce that $\text{Re } f = 0$.

If (f, ψ) is a solution of (5.53) then, writing $\psi = adu_1 \wedge du_2$, applying, respectively, ∂ and ∂^* to the equation we deduce that $\partial^* \partial f = 0$ and $\partial^* \partial a = 0$, that is, f and a are harmonic functions on C_0 . Therefore, f and a each factorize as $r^\lambda G(y)$ (we omit the details but cf. [Jo4, Propn. 2.4]). A function $r^{\lambda-2} G(y)$ is harmonic if and only if G is an eigenfunction of the Laplacian on the link $\mathbb{R}P^3 = \{r = 1\}$ of the singularity of C_0 . The eigenvalue of G is then $\lambda(\lambda - 2)$ and must be non-negative, therefore there are no solutions f, ψ to (5.53) of the form (5.51) with $0 < \lambda < 2$.

The harmonic $(0, 2)$ -form ψ may be written in the uniformizing coordinates as $\psi = a d\bar{u}_1 \wedge d\bar{u}_2$, for some harmonic complex function a . Note that $d\bar{u}_1 \wedge d\bar{u}_2$ is homogeneous of degree -1 in r and so a factorizes as $r^{\lambda-1} G(y)$.

If we assume that both f and a are *homogeneous polynomials* of even degree in $u_1, u_2, \bar{u}_1, \bar{u}_2$ then solving (5.53) with $2 \leq \lambda < 4$ in this case becomes an elementary calculation. We obtain that $\text{Im } f$ must be a constant if $\lambda = 2$ and a harmonic homogeneous quadratic polynomial if $\lambda = 3$. This contributes one real dimension to $d(2)$ and 6 real dimensions to $d(3)$ as a polynomial h can be found to satisfy (5.53). But h then is determined up to an anti-holomorphic homogeneous polynomial of $\bar{u}_1, \bar{u}_2$ of degree $2\lambda - 2$. This contributes further $4\lambda - 2$ real dimensions to $d(\lambda)$. We thus find that $d(1) \geq 2$, $d(2) \geq 7$, and $d(3) \geq 16$. Proposition 5.50, in this context, asserts that these are *all* the solutions of (5.53) for $0 < \lambda \leq 3$ and the above estimates for $d(\lambda)$'s are in fact equalities.

6. PERTURBING THE SINGULAR FIBRES

We can now deduce Theorem A.

Theorem A. *Let M be a compact 7-manifold with a smooth one-parameter family of G_2 -structures given by closed 3-forms $\varphi_T \in \Omega_+^3(M)$, $T > T_0$, defined by the generalized connected sum construction in §2. Let $\tau_T : M \rightarrow S^3$, be a coassociative K3 fibration, with respect to φ_T , defined in Theorem 2.23. Suppose that $\varphi_T + d\eta_T$, is a smooth family of torsion-free G_2 -structures on M , such that $\|\eta_T\|_{L_k^p} < K_{p,k} e^{-\lambda T}$ for each $p > 1$, $k \geq 0$.*

Then there exists T_1 and for any $T > T_1$ and $0 \leq s \leq 1$ a $C^{1\alpha}$ vector field $v_{T,s}$ on M , smooth away from the singular fibres of τ_T and satisfying $\|v_{T,s}\|_{L_k^p} < K_{p,k} s e^{-\lambda T}$ and $\|v_{T,s}\|_{C^1} < K s e^{-\lambda T}$, with support of v contained in a neighbourhood U of the singular fibres of τ_T , and such that $\varphi + s d\eta_T$ vanishes on every singular fibre of the perturbed fibration $\tau_T \circ \exp_{v_{T,s}}^{-1} : M \rightarrow S^3$. The neighbourhood U may be chosen not to meet the neck of M , i.e. $U \subset (W_1(0) \times S^1 \sqcup W_2(0) \times S^1)$.

Theorem A will be deduced by building up on the analysis of the previous two sections. The orbifold fibres of τ_T are parameterized by finitely many disjoint circles. It will suffice to restrict attention to one such S^1 -family. We assume the notation $M_{\text{loc}} = W_{\text{loc}} \times S^1$, $\varphi_{CY} \in \Omega_+^3(M_{\text{loc}})$, $\tau = \tau_T$, $X_0 \subset W_{\text{loc}}$, $X'_0 = X_0 \setminus \{w_0\}$, and the radial coordinate r on X'_0 , as in §4. For each $\theta \in S^1$, define a finite-dimensional space of smooth vector fields

$$\tilde{E}_\theta = \{v_{\mathbf{e},L}|_{X'_0 \times \{\theta\}} : \mathbf{e} \in T_{(w_0, \theta)}(W_{\text{loc}} \times \{\theta\})\},$$

with $v_{\mathbf{e},L}$ as defined in Theorem 4.35.

Theorem 6.54. *Suppose that $1 < k - 4/p < 2$. Then there exists $\varepsilon_0 > 0$ such that for each $\eta \in \Omega^2(M_{\text{loc}})$ with $\|d\eta\|_{C^1(M_{\text{loc}})} < \varepsilon_0$ and $\theta \in S^1$ there is a unique vector field along $X'_0 \times \{\theta\}$, $v(\eta, \theta) \in \tilde{E}_\theta + W_k^p N_{(X'_0 \times \{\theta\})/M_{\text{loc}}}$, with the following properties:*

- (i) $v(0, \theta) = 0$ and v depends smoothly on η and θ ,
- (ii) $\varphi_{CY} + d\eta$ vanishes on $\exp_{v(\eta, \theta)}(X'_0 \times \{\theta\})$.
- (iii) the vector field $v(\eta, \cdot)$ on $X'_0 \times S^1$ is the restriction of some $C^{1, \alpha}$ vector field $v(\eta)$ on M_{loc} smooth away from $X_0 \times S^1$, such that $\|v(\eta)\|_{L_k^p} < C_{p,k} \|d\eta\|_{L_k^p}$

Theorem A is a consequence of Theorem 6.54. Choose T_1 so that the C^1 norm of $d\eta_T$, for $T > T_1$, is smaller than ε_0 given by Theorem 6.54 for each S^1 -family of the singular fibres of τ_T . Further increasing T_1 if necessary, we can ensure that the vector field $v(\eta)$ can be chosen with compact support contained in M_{loc} and with a small C^1 -norm so that $\exp_{v(\eta)}$ is a diffeomorphism. Let U be the union of the neighbourhoods M_{loc} for all S^1 -families of the singular fibres of τ . The vector field $v_{T,s}$ required in Theorem A is then obtained by putting $\eta = s d\eta_T$ and extending the $v(\eta)$ by zero on $M \setminus U$.

In the rest of this section we prove Theorem 6.54.

The proof is based on an application of the Implicit Function Theorem in Banach spaces to a ‘parametric version’ of McLean’s map for the family $X_0 \times S^1$ of coassociative K3 orbifolds

$$F : (v, \eta, \theta) \in (\tilde{E}_0 + W_k^p N_{X'_0/M_{\text{loc}}}) \times \Omega^2(M_{\text{loc}}) \times S^1 \rightarrow \exp_v^*(\varphi_{CY} + R_\theta^* d\eta)|_{X'_0} \in d(\tilde{E}_0 \lrcorner \varphi_{CY}) + (W_{k-1}^p \Omega^3(X'_0) \cap \text{Ker } d), \quad (6.55)$$

where $R_\theta : W_{\text{loc}} \times S^1 \rightarrow W_{\text{loc}} \times S^1$ denotes an isometry given by the translations by θ on the S^1 factor (here we exploited the S^1 -symmetry to make the Banach space of self-dual forms independent of θ). A choice of Banach space for η is not particularly important here as long as it controls e.g. the C^3 norm. We need the following.

Proposition 6.56. *The map F defined in 6.55 is a smooth map of Banach spaces at any (v, η, θ) with sufficiently small $\|v\|$ in $\tilde{E}_0 + W_k^p N_{X'_0/M_{\text{loc}}}$.*

Proof. The map F is linear in η and has an equivariant property

$$F(v, \eta, \theta) = F(v, R_\theta^* \eta, 0). \quad (6.57)$$

Therefore, F is smooth in η, θ and it remains to show that F is smooth in v .

A related argument was carried out by Baier in [Ba, Theorem 2.2.15], see also [JS, Theorem 2.5]. We may disregard a finite-dimensional component $\tilde{E}_0$ and pretend that $v \in W_k^p N_{X'_0/M_{\text{loc}}} \subset C^1 N_{X'_0/M_{\text{loc}}}$, so v has a point-wise estimate $|v| < \varepsilon r$ near w_0 for any $\varepsilon > 0$ (in the metric $g(\varphi_{CY})$). By Proposition 4.29, the Riemannian exponential map is a diffeomorphism between a neighbourhood of the zero section of $N_{X'_0/M_{\text{loc}}}$ and a neighbourhood of X'_0 in M_{loc} of the form $U_\varepsilon = \cup_{x \in X'_0} B_x(\varepsilon r)$ for some $\varepsilon > 0$. Denote by $\tilde{\varphi} \in \Omega^3(U_\varepsilon)$ the pull-back of $\varphi_{CY} + R_\theta^* d\eta$ via this diffeomorphism. Then F , as a function of v , becomes equivalent to

$$v \in W_k^p N_{X'_0/M_{\text{loc}}} \rightarrow v^* \tilde{\varphi} \in W_{k-1}^p \Omega^3(X'_0),$$

for v with sufficiently small in the W_k^p norm. Use a uniformizing coordinate neighbourhood of w_0 in X'_0 and a finite open cover of the compact complement by coordinate neighbourhoods and the respective local trivializations of $N_{X'_0/M_{\text{loc}}}$ to obtain local expressions for v . The pull-back $v^*\tilde{\varphi}$ of a given smooth 3-form is then expressed a cubic polynomial in the partial derivatives of v with coefficients smoothly depending on v . The reader now should have no difficulty to check the smoothness of F using standard results on Sobolev spaces. $\square$

The first partial derivative $D_v F(0, 0, \theta)$ is a composition of the bundle isometry $v \mapsto v \lrcorner \varphi_{CY}$ and the restriction of the exterior derivative operator (4.36) to a complement of its kernel. By Theorem 4.35, $D_v F(0, 0, \theta)$ maps $\tilde{E}_0 \oplus W_k^p N_{X'_0/M_{\text{loc}}}$ isomorphically onto the image of F . Therefore, the implicit function theorem in Banach spaces applies to F and gives for each $\theta_0 \in S^1$ a unique smooth family of fields $v(\eta, \theta)$ in $\tilde{E}_0 \oplus W_k^p N_{X'_0/M_{\text{loc}}}$, for small $\|d\eta\|$ and $|\theta - \theta_0|$, satisfying $v(0, \theta) = 0$ and $F(v(\eta, \theta), \eta, \theta) = 0$. By the compactness of S^1 and the uniqueness of the local solutions $v(\eta, \theta)$, a finite number of these solutions can be patched together to define a vector field $(dR_\theta)_0 v(\eta, \theta)$ on $X'_0 \times S^1$, for all $\|d\eta\| < \varepsilon$, where ε is the smallest of the finitely many upper bounds coming from the local constructions.

This completes the proof of clauses (i) and (ii) of the theorem and it remains to establish the regularity clause (iii).

The interior regularity. We show that the vector fields $v(\eta, \theta)$ defined above are smooth on X'_0 . In general, a $(\varphi_{CY} + d\eta)$ -coassociative submanifold need not be calibrated, so it is not quite sufficient to fall back on the traditional argument for minimal submanifolds. If $v = v_0 + v_1 \in \tilde{E}_0 + W_k^p N_{X'_0/M_{\text{loc}}}$ is a solution of $F(v, \eta, \theta) = 0$ for some η, θ then v_0 is smooth and the equation satisfied by v_1 can be written in the form

$$F_0 + Gv_1 + Q(v_1) = 0. \quad (6.58)$$

Here $Gv_1 = d(v_1 \lrcorner (\varphi_{CY} + d\eta))$ and $F_0 = R_\theta^* d\eta|_{X'_0} + Gv_0$ is a smooth 3-form on X'_0 . It follows from the proof of Proposition 6.56 that the remainder Q is a cubic polynomial in the first derivatives of v_1 with coefficients smoothly depending on v_1 . To deal with this non-linearity note first that $v_1 \in W_k^p N_{X'_0/M_{\text{loc}}}$, with $k - 4/p > 1$, thus in $C^{1,\alpha}$, $\alpha > 0$. Then the first factor in each term $(\nabla v_1)^{\otimes n} \otimes \nabla v_1$ can be considered as a first order linear differential operator with Hölder continuous coefficients acting on the second term. If $\|v_1\|_{C^1}$ is sufficiently small then the sum of the latter linear operator and G is again an overdetermined-elliptic operator, so we still have the standard local regularity estimates [ADN]. As v_1 is already in $C^{1,\alpha}$ we can apply the usual bootstrapping to show that v_1 , and hence also v , is smooth. Then $(dR_\theta)_0(v(\eta, \theta))$ defines a C^∞ vector field on $X'_0 \times S^1$.

Remark. If the 4-form $*_{\varphi_{CY}+d\eta}(\varphi_{CY} + d\eta)$ is closed then $v(\eta, \theta)$ defines a calibrated submanifold and is real analytic. This is a consequence of a general property of minimal submanifolds of a Ricci-flat (hence real analytic [DK]) Riemannian manifold.

$C^{1,\alpha}$ -regularity at $X_0 \times S^1$. The normal bundle $N_{X'_0/M_{\text{loc}}}$ is trivial and W_k^p sections are $C^{1,\alpha}$ and vanish at w_0 together with first derivatives when $k - 4/p > 1$. Multiplying by a smooth cut-off function which is equal to 1 on $X_0 \times S^1$ we obtain an extension of a L_k^p field of normal vectors to a vector field supported on a compact neighbourhood of $X_0 \times S^1$

on M_{loc} . It is clear that such an extension is smooth except at points of $X_0 \times S^1$ where it is $C^{1,\alpha}$. It is easy to see that a $W_k^p N_{X'_0/M_{\text{loc}}}$ section $v(\eta, \theta)$ has an extension to M_{loc} with the same properties as an E_0 component is, by construction, the restriction to $X_0 \times S^1$ of some smooth vector field on M_{loc} .

If $v(\eta)$ has a sufficiently small L_k^p -norm on $X'_0 \times \text{pt}$, then we can achieve estimates $\|v(\eta)\|_{L_k^p(M_{\text{loc}})} < \text{const} \|v(\eta, \cdot)\|_{L_k^p(X'_0 \times S^1)}$, with constants independent of v . The estimates $\|v(\eta, \theta)\|_{p,k} < \text{const} \|d\eta\|_{p,k}$ follow by differentiating the identity $F(\psi(\eta, \theta), \eta, \theta) = 0$ in η, θ and taking account of the symmetry of F in $\theta \in S^1$. If $\|d\eta\|_{C^1}$ is small then $\exp_{v(\eta)}$ is a well-defined diffeomorphism of M_{loc} satisfying the assertions of Theorem 6.54.

7. A NEIGHBOURHOOD OF THE SINGULAR FIBRES

In this section we prove Theorem B and thus complete the proof of the Main Theorem.

Theorem B. *Let M be a compact 7-manifold with a smooth one-parameter family of G_2 -structures given by closed 3-forms $\varphi_T \in \Omega_+^3(M)$, $T > T_1$, defined by the generalized connected sum construction in §2. Let $\tau_T : M \rightarrow S^3$ for $T > T_1$ be a coassociative $K3$ fibration map defined in Theorem 2.23.*

Then there exists $\varepsilon > 0$ so that if $T > T_1$ and an exact form $d\eta \in \Omega^3(M)$ vanishes on every singular fibre of τ_T and $\|d\eta\|_{C^0(M)} < \varepsilon$, relative to the metric $g(\varphi_T)$, then there is a unique smooth vector field $\tilde{v}_T(\eta)$ on M such that:

(i) $\tilde{v}_T$ vanishes on the singular fibres of τ_T and is point-wise orthogonal to each smooth fibre X of τ_T and $\tilde{v}_T \lrcorner \varphi_T|_X$ is L^2 -orthogonal to the harmonic self-dual forms on X relative to the metric $g(\varphi_T)|_X$;

(ii) $\tilde{v}_T(\eta)$ depends smoothly on T and $d\eta$ and $\|\tilde{v}_T\|_{C^1} = O(\|d\eta\|_{C^1})$;

(iii) $\varphi_T + d\eta$ vanishes on the fibres of $\tau_T \circ \exp_{\tilde{v}_T(\eta)}^{-1}$.

The main technical issue in the proof of Theorem B is to establish certain uniform estimates for families of smooth fibres of τ_T in a neighbourhood of the singular fibres. We use the notation of §4: $M_{\text{loc}} = W_{\text{loc}} \times S^1 \subset M$, $\varphi_{CY} \in \Omega_+^3(M_{\text{loc}})$, $X_0 \subset W_{\text{loc}}$, $X'_0 = X_0 \setminus \{w_0\}$, and ‘holomorphic Morse coordinates’ (z_1, z_2, z_3) near w_0 . For $a \in \mathbb{C}$, let $X_{a^2} \subset W_{\text{loc}}$ denote the fibre of τ_T which is expressed near w_0 by the equation $z_1^2 + z_2^2 + z_3^2 = a^2$. Let $X_{a^2, \theta} = X_{a^2} \times \{\theta\} \subset W_{\text{loc}} \times S^1$ and, as before, identify $X_{a^2} = X_{a^2, 0}$, for a marked point $0 \in S^1$. The fibre $X_{a^2, \theta}$ is well-defined for any small $|a|$, say $|a| < \rho$, and is non-singular for $a \neq 0$. Throughout this section any $X_{a^2, \theta}$ is considered with the metric g_{a^2} induced by restriction of $g(\varphi_{CY})$ unless stated otherwise. For each compact smooth coassociative submanifold $X \subset M$, define a decomposition

$$L_k^p(N_{X/M}) = \Gamma^h(N_{X/M}) \oplus \Gamma^\perp(N_{X/M})_{p,k},$$

where

$$\Gamma^h(N_{X/M}) = \{v \in \Gamma(N_{X/M}) \mid (v \lrcorner \varphi_T) \in \mathcal{H}^+(X)\},$$

$$\Gamma^\perp(N_{X/M}) = \{v \in \Gamma(N_{X/M}) \mid (v \lrcorner \varphi_T) \perp_{L^2} \mathcal{H}^+(X)\},$$

and $\Gamma^\perp(N_{X/M})_{p,k}$ means the completion of $\Gamma^\perp(N_{X/M})$ in the L_k^p norm (as before, we require $1 < k - 4/p < 2$). Denote

$$\mathcal{Y} = \{\eta \in \Omega^2(M_{\text{loc}}) \mid d\eta|_{X_{0,\theta}} = 0 \text{ for each } \theta \in S^1\}.$$

The proof of Theorem B builds up on the following.

Proposition 7.59. *There is an $\varepsilon_0 > 0$ so that whenever $\eta \in \mathcal{Y}$ and $\|d\eta\|_{C^1(M_{\text{loc}})} < \varepsilon_0$ the following holds. For each $0 < |a| < \rho$ and $\theta \in S^1$ there is a unique smooth $v = v_{a^2,\theta}(\eta) \in \Gamma^\perp(N_{X_{a^2,\theta}/M_{\text{loc}}})$ such that $v_{a^2,\theta}(0) = 0$, $v_{a^2,\theta}(\eta)$ depends smoothly on η , and $\varphi_{CY} + d\eta$ vanishes on the submanifold $\exp_{v_{a^2,\theta}(\eta)}(X_{a^2,\theta})$.*

For a fixed $\eta \in \mathcal{Y}$ with $\|d\eta\|_{C^1(M_{\text{loc}})} < \varepsilon_0$, the family $v_{a^2,\theta}(\eta)$, for $0 < |a| < \rho$, $\theta \in S^1$ extended by zero over $X_{0,\theta}$, defines a vector field $v(\eta)$ which is C^1 on a neighbourhood M_{loc} of $X_0 \times S^1$ and smooth on $M_{\text{loc}} \setminus X_0 \times S^1$. Furthermore, $\|v(\eta)\|_{C^1(M_{\text{loc}})} < K\|d\eta\|_{C^1(M_{\text{loc}})}$, for some constant K independent of η .

Proof. For each $a \neq 0$, $\theta \in S^1$ we want to obtain $v_{a^2,\theta}(\eta)$ as $v(0, \eta)$, where $v(\beta, \eta)$ is the solution to implicit function problem $F(v(\beta, \eta), \beta, \eta) = 0$ for the extended McLean's map for $X_{a^2,\theta}$

$$F : (v, \beta, \eta) \in \Gamma^\perp(N_{X_{a^2,\theta}/M_{\text{loc}}})_{p,k} \oplus \Gamma^h(N_{X_{a^2,\theta}/M_{\text{loc}}}) \oplus \mathcal{Y} \rightarrow \exp_{v+\beta}^*(\varphi_{CY} + d\eta)|_{X_{a^2,\theta}} \in L_{k-1}^p \Omega^3(X_{a^2,\theta}) \cap \text{Ker } d, \quad (7.60)$$

where $k - 4/p > 1$ we use the completion in the C^3 norm for $\eta \in \mathcal{Y}$. That F is a smooth map between the indicated Banach spaces follows by the argument of Proposition 6.56, up to a change of notation. We have $F(0, 0, 0) = 0$ and the derivative $(D_v F)_{0,0,0} v = d(v \lrcorner \varphi_{CY})|_{X_{a^2,\theta}}$ is an isomorphism of Banach spaces by standard Hodge theory as $H^3(X_{a^2,\theta}, \mathbb{R}) = 0$ for a K3 surface $X_{a^2,\theta}$.

The implicit function theorem in Banach spaces gives a unique family of sections $v(\beta, \eta)$, with $v(0, 0) = 0$, defined for $\max\{|\beta|, \|d\eta\|_{C^1(X_{a^2,\theta})}\} < \delta$ say, but this δ may in general depend on a . (We can choose δ independent of θ by the S^1 -symmetry of φ_{CY} .) To keep track of the relation between constants appearing in the estimates we use.

Proposition 7.61 (Implicit function theorem in Banach spaces). *Suppose that a smooth map $f : E = E_1 \oplus E_2 \rightarrow F$ between Banach spaces has an expansion*

$$f(\xi_1, \xi_2) = (D_1 f(0, 0))\xi_1 + (D_2 f(0, 0))\xi_2 + Q(\xi_1, \xi_2),$$

so that $A = D_1 f(0, 0) : E_1 \rightarrow F$ is an isomorphism of Banach spaces and for $\xi, \zeta \in E$,

$$\|A^{-1}Q(\xi) - A^{-1}Q(\zeta)\| < C(\|\xi\| + \|\zeta\|)\|\xi - \zeta\|,$$

for some constant C . Then there exists a uniquely determined smooth function $\varphi : B_{2,\delta} \rightarrow B_{1,\delta}$, $\varphi(0) = 0$, where $B_{i,\delta} = \{\xi_i \in E_i : \|\xi_i\| < \delta\}$ with $\delta = (4C)^{-1}$, so that all zeros of f in $B_{1,\delta} \times B_{2,\delta}$ are of the form $(\varphi(\xi_2), \xi_2)$.

The proof of Proposition 7.61 is a standard application of the contraction mapping principle.

We require a lower bound on the linearization $(D_1 F)_0$ in v and an upper bound on the quadratic remainder for F so that the constant $(4C)^{-1}$ in Proposition 7.61 for the map F on a manifold X_{a^2} is greater than the norm of $d\eta|_{X_{a^2,\theta}}$ for any sufficiently small $|a| \neq 0$. As noted above, the estimates on $X_{a^2,\theta}$, can be taken independent of $\theta \in S^1$.

We begin with the linear part.

Proposition 7.62. *There exists a $\rho > 0$ such that for any $v \in \Gamma^\perp(N_{X_{a^2,\theta}/M_{\text{loc}}})_{p,k}$ with $0 < |a| < \rho$,*

$$\|v\|_{p,k} < C_{p,k} \|d(v \lrcorner \varphi_{CY})|_{X_{a^2,\theta}}\|_{p,k-1}$$

with a constant $C_{p,k}$ independent of v or a .

Proof. Since the linear map and the norms are symmetric in θ we drop θ from the notation. Consider a sphere $|z| = |a|^{1/2}$ in the coordinate neighbourhood of w_0 in W_{loc} , where $|z|^2 = \sum_{j=1}^3 |z_j|^2$. If $|a|$ is sufficiently small then this sphere intersects X_{a^2} and we can write $X_{a^2} = X_{a^2}^- \cup X_{a^2}^+$, where $X_{a^2}^- = X_{a^2} \setminus \{|z| \leq |a|^{1/2}\}$ and $X_{a^2}^+ = X_{a^2} \cap \{|z| < 2|a|^{1/2}\}$ are two open submanifolds of X_{a^2} . As $a \rightarrow 0$ the metric on $X_{a^2}^-$ is asymptotic in C^∞ to the metric on $X_0 \setminus \{|z| \leq |a|^{1/2}\}$. The other piece $X_{a^2}^+$ with metric rescaled by a constant factor $|a|$ is asymptotic in C^∞ to a complex surface $\Sigma = \{\sum_{j=1}^3 z_j^2 = 1, |z| < |a|\}$ in $\mathbb{C}^3$ with the metric on Σ induced by a Hermitian inner product on $T_{w_0} W_{\text{loc}} \cong \mathbb{C}^3$ defined by the Kähler form $\omega(w_0)$.

The decomposition $X_{a^2} = X_{a^2}^- \cup X_{a^2}^+$ can be thought of as a generalized connected sum $X'_0 \#_{\mathbb{R}P^3} \Sigma$ of two manifolds taken at their ends $\mathbb{R}_{>0} \times \mathbb{R}P^3$. The metric g_{a^2} on the connected sum X_a is C^∞ asymptotic to a metric smoothly interpolating between metrics on compact subsets of X'_0 and Σ .

A family of Riemannian 4-manifolds (X_{a^2}, g_{a^2}) is an instance of the gluing construction studied in [KS, §2]. The estimate that we are interested in is equivalent to a lower bound on the operator

$$d : L_k^p \Omega^+(X_{a^2}) \rightarrow L_{k-1}^p \Omega^3(X_{a^2}). \quad (7.63)$$

The ‘main estimate’ proved in [KS, § 4.1] deals with the invertibility of an elliptic differential operator on a generalized connected sum, as above, when the coefficients of this operator are obtained by gluing the coefficients of Fredholm elliptic operators on the two manifolds with cylindrical ends. The result, in particular, asserts a uniform lower bound on a subspace of finite codimension in the L_k^p domain on a connected sum. Inspection of the proof of this uniform lower bound shows that it remains valid for the overdetermined elliptic operator (7.63) on (X_{a^2}, g_{a^2}) provided that the L^2 kernels of the respective exterior derivatives on self-dual forms on X'_0 and Σ are finite-dimensional. Then the operator (7.63) admits a lower bound, independent of a as $a \rightarrow 0$, on the complement of a finite-dimensional subspace in $L_k^p \Omega^+(X_{a^2})$. The dimension of this space is the sum of dimensions of the spaces of closed self-dual L^2 forms on X'_0 and Σ .

These latter spaces are contained in the L^2 kernels of the Laplacian on self-dual forms on X'_0 and Σ . The dimensions can be computed in the asymptotically cylindrical metrics in the conformal class of X'_0 and Σ as the L^2 norm of 2-forms is conformally invariant

in dimension 4. Recall from Proposition 5.38 that the L^2 kernel is 3-dimensional for $\Omega^+(X'_0)$. The 4-manifold Σ is diffeomorphic to the total space of the tangent bundle TS^2 , so $b^+(\Sigma) = 0$ and the same argument as in Proposition 5.38 shows that the Laplacian on $L_k^p \Omega^+(\Sigma)$ is injective. Thus the uniform lower bound for (7.63) holds on the complement of a 3-dimensional subspace in $\Omega^+(X_{a^2})$. It is *a posteriori* clear that this 3-dimensional space can be taken to be $\mathcal{H}^+(X_{a^2})$. $\square$

The argument of Proposition 6.56 shows that the non-linear remainder Q of the map F consists of polynomial terms in ∇v with coefficients smooth in v . This easily gives an upper bound on Q independent of a, θ for small $|a|$. Therefore, by Propositions 7.61 and 7.62 there is an $\varepsilon > 0$ independent of a, θ so that if $\eta \in \Omega^2(M_{\text{loc}})$ satisfies $\|d\eta|_{X_{a^2, \theta}}\|_{C^1(X_{a^2, \theta})} < \varepsilon$ then the implicit function theorem defines sections $v(0, \eta)$ of $N_{X_{a^2, \theta}/M_{\text{loc}}}$ for every small $|a|$.

The metric on the submanifold $X_{a^2, \theta}$ is independent of $\theta \in S^1$ and in the next result we once again drop θ from the notation. It will be convenient to use Hölder norms commensurable with the L_k^p norms that we required.

Proposition 7.64. *For each X_{a^2} with $a \neq 0$, we have*

$$\|d\eta|_{X_{a^2}}\|_{C^{1, \alpha}(X_{a^2})} < \lambda_{k, \alpha} \|d\eta\|_{C^{1, \alpha}(M_{\text{loc}})} |a|^{1-\alpha}, \quad 0 \leq \alpha < 1,$$

with a constant $\lambda_{k, \alpha}$ independent of η or a .

Estimates similar to that in Proposition 7.64 are proved in [Jo6] and [Lo2]. In the present situation, we have an explicit algebraic local model for the submanifolds X_{a^2} near the singular point of X_0 and some details are simplified.

Proof. Recall from the proof of Proposition 7.62 the decomposition $X_{a^2} = X_{a^2}^- \cup X_{a^2}^+$ as a generalized connected sum and denote by $(d\eta)^\pm$ the restrictions of $d\eta$ to the respective pieces $X_{a^2}^\pm \subset X_{a^2}$. Recall also that $X_{a^2}^-$ is diffeomorphic to $X_0 \setminus \{|z| \leq |a|^{1/2}\}$. As $X'_0 = X_0 \setminus \{|z| \leq |a|^{1/2}\}$ is compact and non-singular, its injectivity radius is bounded away from zero, furthermore, it is not difficult to check that the injectivity radius is $O(|a|)$, as $|a| \rightarrow 0$. We deduce that for $|a|$ sufficiently small, each $X_{a^2}^-$ is a well-defined deformation of X'_0 defined by a section ν_{a^2} of the normal bundle of the latter submanifold, using the exponential map for $g(\varphi_{CY})$. The $\nu_{a^2} \rightarrow 0$ depends smoothly on a and vanishes when $a = 0$. As $d\eta$ is smooth and vanishes on X'_0 and the metric on $X_{a^2}^-$ converges in C^∞ to the metric on X'_0 , we find that the C^k norm of $(d\eta)^-$ is a smooth function of a . This function vanishes at $a = 0$, hence is bounded by a constant multiple of $\|d\eta\|_{C^k(M_{\text{loc}})} |a|$.

The form $(d\eta)^+$ is defined on a coordinate neighbourhood $\{|z| < |a|^{1/2}\} \times S^1$ of w_0 in M_{loc} . We may assume, by rescaling $z \in \mathbb{C}^3$ if necessary, that the matrix of the inner product defined by $g(\varphi_{CY})$ on $T_{w_0}W_{\text{loc}}$ has determinant 1. The metric $g(\varphi_{CY})$ on the coordinate neighbourhood may be written as $g_0 + O(|z|)$, where g_0 denotes the value of $g(\varphi_{CY})$ at w_0 and $O(|z|)$ is estimated independently of θ . A similar local expansion is valid for the derivatives of $g(\varphi_{CY})$.

The parameterizations of local submanifolds $X_{a^2}^+$ may be taken to be homogeneous of order 1 in a . The coefficients of the k -th derivatives of induced metric $g(X_{a^2}^+)$ on $X_{a^2}^+$

then have a local extension with the leading term homogeneous of order $2 - k$ in a and a remainder $O(|a|^{3-k})$ as $a \rightarrow 0$ ($d\theta$ vanishes on each X_{a^2}). The difference between the restriction of the Levi-Civita connection of $g(\varphi_{CY})$ on the ambient M_{loc} to $X_{a^2}^+$ and the Levi-Civita connection of the metric $g(X_{a^2}^+)$ induced by $g(\varphi_{CY})$ on X_{a^2} is determined by second fundamental form of $g(X_{a^2}^+)$. Recall that for a submanifold defined by submersion τ_T the second fundamental form is the quotient of the Hessian of τ_T and the gradient of τ_T . Thus its point-wise norm has leading term homogeneous of order a^{-1} , as measured by $g(X_{a^2}^+)$. As $d\eta$ vanishes on X_0 , this 2-form vanishes on $T_{w_0}W_{\text{loc}}$, the Zariski tangent space of X_0 at w_0 , so the coefficients of $d\eta|_{W_{\text{loc}}}$ have zeros of order two at w_0 and the covariant derivative has zero of order one. It follows that for $0 < |a| < \rho$ with sufficiently small ρ , the $C^{1,\alpha}$ norm of $d\eta|_{X_{a^2}^+}$ measured using the Levi-Civita of $g(X_{a^2}^+)$ is $O(\|d\eta\|_{C^k(M_{\text{loc}})}|a|^{1-\alpha})$.

As $\|d\eta|_{X_{a^2}}\|_{C^k(X_{a^2})} \leq \max\{\|d\eta^-|_{X_{a^2}^-}\|_{C^k(X_{a^2}^-)}, \|d\eta^+|_{X_{a^2}^+}\|_{C^k(X_{a^2}^+)}\}$ the proposition is proved. $\square$

The local deformation $\exp_{v(\beta,\eta)}(X_{a^2,\theta})$ is well-defined if $v(\beta,\eta)$ is small in the uniform norm so that the tubular neighbourhood theorem is applicable to $X_{a^2,\theta}$. It is not difficult to see that, for any small a , a suitable neighbourhood of the zero section of $N_{X_{a^2,\theta}/M_{\text{loc}}}$ is $\|v\|_{C^0} < \varepsilon_1 |a|$, for a sufficiently small constant $\varepsilon_1 > 0$ independent of a .

For a small $d\eta$ we can estimate on $X_{a^2,\theta}$, $\|v(0,\eta)\| = O(\|(D_2v)_{0,0}\| \|d\eta|_{X_{a^2,\theta}}\|)$. As $(D_2v)_{0,0} = (D_1F)_{0,0,0}^{-1}(D_3F)_{0,0,0}$ this gives

$$\|v(0,\eta)\|_{C^1(X_{a^2,\theta})} = O(|a|), \quad a \rightarrow 0, \quad (7.65)$$

using Proposition 7.62 and Lemma 7.64. In particular, for a sufficiently small $d\eta$ the deformations $\exp_{v(0,\eta)}(X_{a^2,\theta})$ are well-defined for every small $|a|$.

The regularity results for the coassociative submanifolds [HL] imply that $v(\beta,\eta)$ is a smooth section over $X_{a^2,\theta}$, as $v(\beta,\eta)$ is already in C^1 . By construction, $\exp_{v(\beta,\eta)}(X_{a^2,\theta})$ coincides with $\exp_{v(0,\eta)}(\exp_{v(\beta,0)}(X_{a^2,\theta}))$. Therefore, the local families $v(\beta,\eta)$ can be patched together to define a smooth vector field $v(\eta)$ on $(\cup_{0 < |a| < \rho} X_{a^2}) \times S^1$, for some $\rho > 0$. This vector field has a Lipschitz continuous extension by zero over $\cup_{\theta \in S^1} X_{0,\theta}$.

Differentiating the identity $F(v(\beta,\eta), \beta, \eta) = 0$ in β at $\beta = 0$ we find that the derivative of $v(\eta)$ in the directions orthogonal to the fibres $X_{a^2,\theta}$ satisfies

$$(D_1F)_{v(0,\eta),0,\eta}(D_1v)_{0,\eta} + (D_2F)_{v(0,\eta),0,\eta} = 0.$$

Using explicit expressions for the derivatives of F and the vanishing of φ_{CY} on $X_{a^2,\theta}$ we obtain

$$d(((D_1v)_{0,\eta}\beta) \lrcorner \varphi_{CY})|_{X_{a^2,\theta}} = -d(\beta \lrcorner d\eta)|_{X_{a^2,\theta}}.$$

The operator $d(\cdot \lrcorner \varphi_{CY})|_{X_{a^2,\theta}}$ is bounded below independent of a by Proposition 7.62 and we deduce that $\|(D_1v)_{0,\eta}\|_{C^0(X_{a^2,\theta})} < K\|d\eta\|_{C^1(M_{\text{loc}})}$, $a \neq 0$.

We adapt the same method to show that $v(\eta)$ is continuously differentiable at any point in $X_0 \times S^1$. Differentiating the identity $F(v(\beta,\eta), \beta, \eta) = 0$ in β at $\beta = 0$ we find that the

derivative of $v(\eta)$ in the directions orthogonal to the fibres $X_{a^2, \theta}$ satisfies

$$(D_1 F)_{v(0, \eta), 0, \eta} (D_1 v)_{0, \eta} + (D_2 F)_{v(0, \eta), 0, \eta} = 0.$$

The self-dual harmonic forms on $X_{a^2, \theta}$ are spanned by the Kähler form $\omega|_{X_{a^2, \theta}}$ and the real and imaginary parts of the $(2, 0)$ -form $(\partial\tau)^\sharp \lrcorner \Omega$ induced by the Calabi–Yau structure (ω, Ω) on $W_{\text{loc}} \times \{\theta\}$. The definition of $\Gamma^h(N_{X_{a^2, \theta}/M})$ therefore can be extended to the smooth subset $X'_{0, \theta} \subset X_{0, \theta}$. The resulting $\Gamma^h(N_{X'_{0, \theta}/M})$ can be thought of as a limit of $\Gamma^h(N_{X_{a^2, \theta}/M})$ as $a \rightarrow 0$, in the sense that the 4-manifold $X'_{0, \theta}$ is diffeomorphic to an open subset of $X_{a^2, \theta}$ and the metric on $X'_{0, \theta}$ is a C^∞ limit of the metrics on this subset. Using explicit expressions for the derivatives of F and taking the limit as $a \rightarrow 0$ we obtain

$$d(((D_1 v)_{0, \eta} \beta) \lrcorner \varphi_{CY})|_{X'_{0, \theta}} = -d(\beta \lrcorner d\eta)|_{X'_{0, \theta}}$$

The operator $d(\cdot \lrcorner \varphi_{CY})|_{X'_{0, \theta}}$ is injective on $\Gamma^\perp(N_{X'_{0, \theta}/M_{\text{loc}}})$. Applying a left inverse we determine $(D_1 v)_{0, \eta}$ at any point in $X'_{0, \theta}$. This shows the continuity of $(D_1 v)_{0, \eta}$ away from critical point of τ_T . The estimate (7.65) shows that the derivative of $v(\eta)$ is Lipschitz continuous at each point in $w_0 \times S^1$, thus $v(\eta)$ is C^1 -regular on all of M_{loc} . The higher order derivatives of $v(\eta)$ can be handled by a similar method but the expressions become cumbersome; we omit the details.

We estimated the ‘intrinsic’ first derivatives of $v(\eta)$ on $X_{a^2, \theta}$ and in the transverse directions by $\|d\eta\|_{C^1(M_{\text{loc}})}$. To obtain the desired estimate of the C^1 norm of $v(\eta)$ on M_{loc} we recall that the second fundamental form of $X_{a^2, \theta}$ is $O(|a|^{-1})$, as $a \rightarrow 0$, near the critical points of τ_T and bounded on the complement of a neighbourhood of the critical points. It follows that $\|v\|_{C^1(M_{\text{loc}})} = O(\|d\eta\|_{C^1(M_{\text{loc}})})$. This completes the proof of Proposition 7.59. $\square$

The singular fibres of τ_T occur in finitely many S^1 -families. Applying Proposition 7.59 to each S^1 -family we obtain for any $\eta \in \Omega^2(M_{\text{loc}})$ with small $\|d\eta\|_{C^1}$ a vector field $\tilde{v}_T(\eta)$ satisfying the assertions of Theorem B except that $\tilde{v}_T(\eta)$ is only defined on a neighbourhood U of the singular fibres of τ_T .

The implicit function argument for the extended McLean’s map $F(v, \beta, \eta)$ as in (7.60) for a fibre X of τ_T in $M \setminus U$ gives a uniquely determined smooth local family $v(\beta, \eta)$ of sections in $\Gamma^\perp(N_{X/M})$, so that $v(0, 0) = 0$ and $\varphi_T + d\eta$ vanishes on $\exp_{v(\beta, \eta)}(X)$.

The estimates on $v(\beta, \eta)$ are similar to those in Proposition 7.59 but easier as the curvature of the fibres in $M \setminus U$ is bounded independent of the fibre. The complement $M \setminus U$ is a family of non-singular fibres of τ_T . Recall from §2 that the metric on each of these fibres is up to a small deformation a metric in a *compact* closure of the finite-dimensional family of metrics on the fibres of holomorphic fibrations $\tau^{(j)}$, $j = 1, 2$, restricted to the asymptotically cylindrical ends of W_j . The deformation is bounded, independent of T or the fibre, in C^k norm for each k . It follows that the lower bound on $(D_1 F)_{0, 0, 0}$ and the upper bound on the quadratic remainder can be taken independent of $X \subset M \setminus U$. Then the sections $v(\beta, \eta)$ are defined for $\|d\eta\|_{C^1(M)} < \varepsilon$ where ε can be taken independent of T or the fibre X .

Using once again the property that for a fixed β the sections $v(\beta, \eta)$ define deformations of $\exp_{v(\beta, 0)}(X)$, a fibre near X , we obtain that $v(\beta, \eta)$ induces a vector field satisfying the assertions of Theorem *B* except that it is only defined on an open neighbourhood of X in M . The vector fields obtained from $v(\beta, \eta)$ for different X agree on the overlaps of their domains by the uniqueness part of the implicit function theorem. As $M \setminus U$ is compact an extension of $v(\eta)$ from U to M is obtained by patching with a finite number of these local vector fields.

REFERENCES

- [Ac] B. S. Acharya. *Exceptional mirror symmetry*. In Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), pages 1–14. Amer. Math. Soc., 2001.
- [ADN] S. Agmon, A. Douglis, and L. Nirenberg. *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II*. Comm. Pure Appl. Math. **17** (1964), 35–92.
- [AF] A. Andreotti and T. Frankel. *The second Lefschetz theorem on hyperplane sections*. In Global Analysis (Papers in Honor of K. Kodaira), pages 1–20. Univ. Tokyo Press, Tokyo, 1969.
- [As] P.S. Aspinwall. *K3 surfaces and string duality*. Fields, strings and duality (Boulder, CO, 1996), 421–540, World Sci. Publ., River Edge, NJ, 1997. arXiv:hep-th/9611137
- [APS] M.F. Atiyah, V.K. Patodi, and I.M. Singer. *Spectral asymmetry and Riemannian geometry, I*. Math. Proc. Camb. Phil. Soc. **77** (1975), 97–118.
- [Ba] P.D. Baier. *Special Lagrangian geometry*. D.Phil thesis, University of Oxford, 2001.
- [B] W.L. Baily. *The decomposition theorem for V-manifolds*. Amer. J. Math **78** (1956), 862–888.
- [Bi] O. Biquard. *Fibrés paraboliques stables et connexions singulières plates*. Bull. Soc. Math. France **119** (1991), 231–257.
- [Br] R.L. Bryant. *Calibrated embeddings in the special Lagrangian and coassociative cases*. Ann. Global Anal. Geom. **18** (2000), 405–435.
- [DK] D.M. DeTurck and J.L. Kazdan. *Some regularity theorems in Riemannian geometry*. Ann. Sci. École Norm. Sup. (4) **14** (1981), 249–260.
- [FM] R. Friedman and J.W. Morgan. *Smooth four-manifolds and complex surfaces*. Springer-Verlag, Berlin, 1994.
- [Go] E. Goldstein. *Calibrated fibrations*. Comm. Anal. Geom. **10** (2002), 127–150.
- [GYZ] S. Gukov, S.-T. Yau, and E. Zaslow. *Duality and fibrations on G_2 manifolds*. Turkish J. Math. **27** (2003), 61–97.
- [HL] R. Harvey and H.B. Lawson, Jr. *Calibrated geometries*. Acta Math. **148** (1982), 47–157.
- [Ha] B. Hassett. *Potential density of rational points on algebraic varieties*. In Higher dimensional varieties and rational points (Budapest, 2001), vol. 12 of Bolyai Soc. Math. Stud., pages 223–282. Springer, 2003.
- [Is] V.A. Iskovskih. *Fano threefolds. I*. Izv. Akad. Nauk SSSR Ser. Mat. **41** (1977), 516–562, 717. English translation: Math. USSR-Izvestia **11** (1977) 485–527.
- [Jo1] D.D. Joyce. *Compact Riemannian 7-manifolds with holonomy G_2 . I, II*. J. Diff. Geom. **43** (1996), 291–328, 329–375.
- [Jo2] D.D. Joyce. *Compact manifolds with special holonomy*. OUP Mathematical Monographs series, 2000.
- [Jo3] D.D. Joyce. *Singularities of special Lagrangian fibrations and the SYZ conjecture*. Comm. Anal. Geom. **11** (2003), 859–907.
- [Jo4] D.D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. I. Regularity*. Ann. Global Anal. Geom. **25** (2004), 201–251.

- [Jo6] D.D. Joyce. *Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case*. Ann. Global Anal. Geom. **26** (2004), 1–58.
- [Jo7] D.D. Joyce. *Riemannian holonomy groups and calibrated geometry*. Oxford University Press, 2007.
- [JS] D.D. Joyce and S. Salur. *Deformations of asymptotically cylindrical coassociative submanifolds with fixed boundary*. Geom. Topol. **9** (2005), 1115–1146 (electronic).
- [Ko] A.G. Kovalev. *Twisted connected sums and special Riemannian holonomy*. J. Reine Angew. Math. **565** (2003), 125–160.
- [KLe] A.G. Kovalev and Nam-Hoon Lee. *K3 surfaces with non-symplectic involution and compact irreducible G_2 -manifolds*. Preprint arXiv:0810.0957 version 3, September 2009.
- [KN] A.G. Kovalev and J. Nordström. *Asymptotically cylindrical 7-manifolds of holonomy G_2 with applications to compact irreducible G_2 -manifolds*. Preprint arXiv:0907.0497 version 3, August 2009.
- [KS] A.G. Kovalev and M.A. Singer. *Gluing theorems for complete anti-self-dual spaces*. Geom. Funct. Anal. **11** (2001), 1229–1281.
- [La] S. Lang. *Differential manifolds*. Addison-Wesley, 1972.
- [LM] R.B. Lockhart and R.C. McOwen. *Elliptic differential operators on noncompact manifolds*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. **12** (1985), 409–447.
- [Lo1] J.D. Lotay. *Coassociative 4-folds with conical singularities*. Comm. Anal. Geom. **15** (2007), 891–946.
- [Lo2] J.D. Lotay. *Desingularization of coassociative 4-folds with conical singularities*. Geom. Funct. Anal. **18** (2009), 2055–2100.
- [Lo3] J.D. Lotay. *Examples of coassociative 4-folds with conical singularities*. preprint arXiv:0910.5092, October 2009.
- [McL] R.C. McLean. *Deformations of calibrated submanifolds*. Comm. Anal. Geom. **6** (1998), 705–747.
- [MP] V.G. Maz’ja and B.A. Plamenevski. *Estimates on L^p and Hölder classes on the Miranda–Agmon maximum principle for solutions of elliptic boundary value problems with singular points on the boundary*. Math. Nachr. **81** (1978), 25–82. (Russian) English translation: Amer. Math. Soc. Transl. Ser. 2 **123** (1984) 1–56.
- [Me] R.B. Melrose. *The Atiyah–Patodi–Singer index theorem*. A K Peters Ltd., Wellesley, MA, 1993.
- [MMR] J.W. Morgan, T. Mrowka, and D. Ruberman. *The L^2 -moduli space and a vanishing theorem for Donaldson polynomial invariants*. International Press, Boston, 1994.
- [N] J. Nordström. *Deformations of glued G_2 -manifolds*. Comm. Anal. Geom. **17** (2009), 481–503. arXiv:0809.4055
- [Sa] S.M. Salamon. *Riemannian geometry and holonomy groups*. Pitman Res. Notes in Math. 201. Longman, Harlow, 1989.
- [Sh] V.V. Shokurov. *Smoothness of a general anticanonical divisor on a Fano variety*. Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), 430–441. English translation: Math. USSR-Izvestia **14** (1980) 395–405.
- [SYZ] A. Strominger, S.-T. Yau, and E. Zaslow. *Mirror Symmetry is T-Duality*. Nucl. Phys. B **479** (1996), 243–259.
- [TY] G. Tian and S.-T. Yau. *Complete Kähler manifolds with zero Ricci curvature. I*. J. Amer. Math. Soc. **3** (1990), 579–609.
- [Y] S.-T. Yau. *On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation. I*. Comm. Pure Appl. Math., **31** (1978), 339–411.

DPMMS, UNIVERSITY OF CAMBRIDGE, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, GREAT BRITAIN
E-mail address: a.g.kovalev@dpmms.cam.ac.uk